

Fermat Primes.

César Aguilera.

August 4, 2025

Abstract

Properties related to Fermat Primes, their connectedness with Mersenne Numbers and the reason why there are not infinite many Mersenne Primes and Perfect Numbers.

Introduction.

The Fermat Primes, discovered by the french Mathematician Pierre de Fermat, are prime numbers of the form $2^{2^n} + 1$ for $n \geq 0$, there are only five known numbers of this type (3,5,17,257,65537).

Theorem. (1)

Given the matrix $A = \begin{vmatrix} n & 1 \\ n+1 & 2n+1 \end{vmatrix}$ with Determinant $|A|$, then:

$$A = \begin{vmatrix} n & 1 \\ n+1 & 2n+1 \end{vmatrix} = \begin{vmatrix} n & 1 \\ n+1 & F_k \end{vmatrix} \Rightarrow \begin{vmatrix} |A|+1 & 1 \\ |A|+2 & F_{(k+1)} \end{vmatrix} \quad (1)$$

This is:

$$\begin{vmatrix} 1 & 1 \\ 2 & 3 \end{vmatrix} = 1 \Rightarrow \begin{vmatrix} 2 & 1 \\ 3 & 5 \end{vmatrix} = 7 \Rightarrow \begin{vmatrix} 8 & 1 \\ 9 & 17 \end{vmatrix} = 127 \Rightarrow \begin{vmatrix} 128 & 1 \\ 129 & 257 \end{vmatrix} = 32767 \Rightarrow \begin{vmatrix} 32768 & 1 \\ 32769 & 65537 \end{vmatrix}$$

The matrix $\mathbf{A}$ is a Fibonacci Box,[1] and they have the next properties:

$$A = \underbrace{\begin{bmatrix} x & w \\ y & z \end{bmatrix}}_{[a,b,c]} \Rightarrow \begin{aligned} y &= x + w \\ z &= y + x \\ a &= (z)(w) \\ b &= (2)(x)(y) \\ c &= (z)(x) + (y)(w) \end{aligned}$$

so we take $z = 2^{2^{(n-1)}} + 1 = F_k$ and $w = 1$ and then we can find the other terms x and y .

From matrix A we have $\begin{vmatrix} 2^{(2^{(n-1)}-1)} & 1 \\ 2^{(2^{(n-1)}-1)} + 1 & 2^{2^{(n-1)}} + 1 \end{vmatrix}$ where $|A| = 2^{(2^n-1)} - 1$

thus, we can find the Primitive Pythagorean Triple [a,b,c]:

$$a = (z)(w) = (2^{2^{(n-1)}} + 1)(1) = 2^{2^{(n-1)}} + 1 = F_k$$

$$b = (2)(x)(y) = (2)(2^{(2^{(n-1)}-1)})(2^{(2^{(n-1)}-1)} + 1) = 2^{2^{n-1}} + 2^{2^n-1}$$

$$c = (z)(x) + (y)(w) \\ = (2^{2^{(n-1)}} + 1)(2^{(2^{(n-1)}-1)}) + (2^{(2^{(n-1)}-1)} + 1)(1) = 2^{2^{n-1}} + 2^{2^n-1} + 1$$

So, if:

$$\text{Fermat primes } F_k = 2^{2^{(n-1)}} + 1 = \{3, 5, 17, 257, 65537\}$$

and

$$|A| = 2^{(2^n-1)} - 1 = \{0, 1, 7, 127, 32767\} \text{ then, from (1): } 2|A| + 3 = F_{(k+1)}$$


Also if

$$|A|_k = n_k \rightarrow |A|_{(k+1)} = ((F_{(k+1)})(|A|_k + 1)) - (|A|_k + 2)(1) \\ = 2(|A|_k)^2 + 4|A|_k + 1 \text{ or } n_{(k+1)} = 2(n_k)^2 + 4n_k + 1. \quad (2)$$

In other words, if the polynomial $2n^2 + 4n + 1$ takes the values of $n = |A_k|$, we obtain $|A_{k+1}|$.

Primitive Pythagorean Triples related to Areas,

Graphic representation of $\mathbf{A} = \begin{vmatrix} n & 1 \\ n+1 & 2n+1 \end{vmatrix}$ as a determinant.



Areas of segments.

$$A = F = x_2 \cdot y_1 = n + 1$$

$$B = E = \frac{x_1 \cdot y_1}{2} = \frac{n \cdot (n+1)}{2} = \frac{n^2 + n}{2}$$

$$C = D = \frac{x_2 \cdot y_2}{2} = \frac{y_2}{2} = \frac{2n+1}{2}$$

Figure 1: Determinant.

[2]

As the matrix is a Fibonacci box, we calculate the Primitive Pythagorean Triple.

$$a = \{2C \vee 2D\} = 2 \cdot \frac{2n+1}{2} = 2n+1 = F_k = C + D.$$

$$b = \{4B \vee 4E\} = 4 \cdot \frac{n^2+n}{2} = 2n^2 + 2n$$

$$c = \{4B \vee 4E\} + 1 = 4 \cdot \frac{n^2+n}{2} + 1 = 2n^2 + 2n + 1$$

So, the Pythagorean Triple related to Areas is: $\underbrace{[2n+1]}_a, \underbrace{[2n^2+2n]}_b, \underbrace{[2n^2+2n+1]}_c$

now, we can perform Pythagorean Triples whose shortest leg is the n-th Fermat number, we can see them in the next table with only Fermat Primes.

Table 1

$a = F_k = 2n + 1$	$b = 2n^2 + 2n$	$2n^2 + 2n + 1$
3	4	5
5	12	13
17	144	145
257	33024	33025
65537	2147549184	2147549185

note that:

$$\sqrt{(2n^2 + 2n) + (2n^2 + 2n + 1)} = \sqrt{(2n^2 + 2n + 1)^2 - (2n^2 + 2n)^2} = 2n + 1$$

this is the short leg F_k

To obtain these Pythagorean Triples, we need to transform from:

$$2^{2^{(n-1)}} + 1 \text{ to the short leg } 2n + 1 = a$$

$$2^{2^n} + 2^{2^{n+1}-1} \text{ to the long leg } 2n^2 + 2n = b.$$

$$2^{2^n} + 2^{2^{n+1}-1} + 1 \text{ to the hypotenuse } 2n^2 + 2n + 1 = c.$$

As these Pythagorean Triples are related to areas, we can represent them this way

$$a = 2n + 1 = F_k, b = \text{sqr}t \left[\int_1^{F_k} (x^3 - x) dx \right], c = \text{sqr}t \left[\int_1^{F_k} (x^3 - x) dx + 1 \right]$$

[3]

Properties related to the Dot product $|\vec{a} \cdot \vec{b}|$.

Given two vectors $\mathbf{a}$ and $\mathbf{b}$, such that:

$$\mathbf{a} = (2^{(2^{(n-1)}-1)}, 2^{(2^{(n-1)}-1)+1}) = (n, n+1) \text{ and } \mathbf{b} = (1, 2^{2^{(n-1)}}+1) = (1, 2n+1) \text{ then } \frac{|\vec{a} \cdot \vec{b}| - |A|}{2} = F_k,$$

this is, the dot product of $\underbrace{\begin{bmatrix} n \\ n+1 \end{bmatrix}}_{\vec{a}} \cdot \underbrace{\begin{bmatrix} 1 \\ 2n+1 \end{bmatrix}}_{\vec{b}}$ minus $|A|$, and all that divided by 2

is a Fermat Number, so we have:

$$|\vec{a} \cdot \vec{b}| = 2n^2 + 4n + 1, |A| = 2n^2 - 1$$

$$|\vec{a} \cdot \vec{b}| - |A| = 4n + 2 \Rightarrow \frac{4n+2}{2} = 2n + 1.$$

The dot product $|\vec{a} \cdot \vec{b}|$ is also (2) and have the same value of a Determinant so,

$$\frac{|\vec{a} \cdot \vec{b}| - |A|}{2} = F_k \text{ turns to } \frac{|A|_k - |A|_{k-1}}{2} = F_k \text{ this is:}$$

$$\frac{2n^2 + 4n + 1 - (2(n-1)^2 + 4(n-1) + 1)}{2} = \frac{4n+2}{2} = 2n + 1 = F_k$$

Note that $2n + 1$ which represents the Fermat Numbers becomes the *short leg* of the pythagorean triangle (a, b, c) .

$$\text{another property is: } a + b = \underbrace{2n + 1}_a + \underbrace{2n^2 + 2n}_b = 2n^2 + 4n + 1 \text{ thus } a + b = |\vec{a} \cdot \vec{b}|$$

Also note that:

$$(2n^2 + 2n + 1)(4n^2 + 4n + 2) - (2n^2 + 4n + 1)^2 = (2n^2 - 1)^2$$

this is:

$$(|\vec{a}| |\vec{b}|)^2 - |\vec{a} \cdot \vec{b}|^2 = |A|^2$$

from (1) we have:

$$A_{k+1} = \begin{vmatrix} \sqrt{(|\vec{a}| |\vec{b}|)^2 - |\vec{a} \cdot \vec{b}|^2} + 1 & 1 \\ \sqrt{(|\vec{a}| |\vec{b}|)^2 - |\vec{a} \cdot \vec{b}|^2} + 2 & 2\sqrt{(|\vec{a}| |\vec{b}|)^2 - |\vec{a} \cdot \vec{b}|^2} + 3 \end{vmatrix}$$

$$\text{Since } \left\{ \prod_1^k F_k \right\} + 2 = F_{k_n} \Rightarrow 3 \left\{ \prod_1^k 2\sqrt{(|\vec{a}_k| |\vec{b}_k|)^2 - |\vec{a}_k \cdot \vec{b}_k|^2} + 3 \right\} + 2 = F_{k_n}$$

example:

$$\begin{aligned} & 3(2\sqrt{(\sqrt{5}\sqrt{10})^2 - 7^2} + 3) \times (2\sqrt{(\sqrt{13}\sqrt{26})^2 - 17^2} + 3) \\ & \times (2\sqrt{(\sqrt{145}\sqrt{290})^2 - 161^2} + 3) \times (2\sqrt{(\sqrt{33025}\sqrt{66050})^2 - 33281^2} + 3) + 2 \\ & = 4294967297 \end{aligned}$$

Euler-Legendré-Gauss Theorems.

if every Mersenne number $M_n \equiv 2^n - 1$ we can write:

$$|\vec{a}_k| = \sqrt{\left(2^{\frac{M_{n_k}-1}{2}}\right)^2 + \left(2^{\frac{M_{n_k}-1}{2}} + 1\right)^2}$$

$$|\vec{b}_k| = \sqrt{2} \sqrt{\left(2^{\frac{M_{n_k}-1}{2}}\right)^2 + \left(2^{\frac{M_{n_k}-1}{2}} + 1\right)^2}$$

$$(d_k)^2 = \left(2^{\frac{M_{n_k}-1}{2}}\right)^2 + \left(2^{\frac{M_{n_k}-1}{2}} - 1\right)^2$$

so, from the *Euler-Legendré-Gauss* Quadratic Reciprocity Theorem:

if $(d_k)^2$ is prime then $(d_k)^2 \equiv 1 \pmod{4}$

$$\text{since } (d_k)^2 = |A_{k-1}|^2 + |A_{k-1} + 1|^2$$

$$\text{and } \left(|\vec{a}_k \cdot \vec{b}_k| - |\vec{a}_k|^2\right)^2 = |\vec{a}_{k+1} \cdot \vec{b}_{k+1}| - |\vec{a}_{k+1}|^2$$

from *Legendrés Sum of Three Squares Theorem*:

$$|\vec{a}_k \cdot \vec{b}_k| - (d_k)^2 = \left(|\vec{a}_{k-1} \cdot \vec{b}_{k-1}| - |\vec{a}_{k-1}|^2\right)^2 + \left(|\vec{a}_{k-1} \cdot \vec{b}_{k-1}| - |\vec{a}_{k-1}|^2\right)^2 + \left(|\vec{a}_{k-1} \cdot \vec{b}_{k-1}| - |\vec{a}_{k-1}|^2\right)^2$$

or

$$|\vec{a}_k \cdot \vec{b}_k| - (d_k)^2 = (F_{k-1} - 1)^2 + (F_{k-1} - 1)^2 + (F_{k-1} - 1)^2$$

Given $\begin{vmatrix} n & 1 \\ n+1 & F_k \end{vmatrix} = |A|_k \Rightarrow (d_k)^2 = |A|_k - F_k + 3$ and $\frac{F_k-3}{2} = |A|_{k-1}$
 $\Rightarrow 2|A|_{k-1} + 3 = F_k$, since $(d_k)^2 \equiv 1 \pmod{4}$
the recurrent division of $\frac{(d_k)^2-1}{4} = |A|_{k-1}$

Example:

$$A = \begin{vmatrix} 32768 & 1 \\ 32769 & 65537_{F_k} \end{vmatrix} \Rightarrow |A| = \underbrace{2147483647}_{\text{Mersenne prime and } n-1}$$

$$\frac{(d^2)-1}{4} = \frac{2147418113-1}{4} = 536854528 \dots \frac{131068}{4} = 32767$$

$$A = \begin{vmatrix} 128 & 1 \\ 129 & 257_{F_k} \end{vmatrix} \Rightarrow |A| = 32767$$

$$\frac{(d^2)-1}{4} = \frac{32513-1}{4} = \underbrace{8128}_{\text{Perfect number}} \dots \frac{508}{4} = \underbrace{127}_{\text{Mersenne prime and } n-1}$$

$$A = \begin{vmatrix} 8 & 1 \\ 9 & 17_{F_k} \end{vmatrix} \Rightarrow |A| = 127$$

$$\frac{(d^2)-1}{4} = \frac{113-1}{4} = \underbrace{28}_{\text{Perfect number}} \dots \frac{28}{4} = \underbrace{7}_{\text{Mersenne prime and } n-1}$$

$$A = \begin{vmatrix} 2 & 1 \\ 3 & 5_{F_k} \end{vmatrix} \Rightarrow |A| = 7$$

$$\frac{(d^2)-1}{4} = \frac{5-1}{4} = \frac{4}{4} = 1 \Rightarrow A = \begin{vmatrix} 1 & 1 \\ 2 & 3_{F_k} \end{vmatrix} \Rightarrow |A| = 1$$

while we divide, notice the presence of **Perfect Numbers** and almost **Perfect Numbers** as well **Mersenne Primes** and almost **Mersenne Primes**.
In the matrix of the first non prime Fermat Number we have:

$$A = \begin{vmatrix} 2147483648 & 1 \\ 2147483649 & 4294967297_{F_k} \end{vmatrix} \Rightarrow |A| = 9223372036854775807$$

$$\frac{(d^2)-1}{4} = \frac{9223372032559808512}{4} = \underbrace{2305843008139952128}_{\text{Perfect Number}} \text{ then}$$

$$\underbrace{2305843009213693951}_{\text{Mersenne prime}} - \underbrace{2305843008139952128}_{\text{Perfect Number}} = \underbrace{1073741823}_{n-1} \text{ and then}$$

$$A = \begin{vmatrix} 1073741824 & 1 \\ 1073741825 & 2147483649 \end{vmatrix} \Rightarrow |A| = \underbrace{2305843009213693951}_{\text{Mersenne prime}}$$

The $\underbrace{2147483647}_{\text{Mersenne prime}}$ corresponds to $\underbrace{2305843008139952128}_{\text{Perfect Number}}$

Also:

$$A = \begin{vmatrix} 128 & 1 \\ 129 & 257_{F_k} \end{vmatrix} \Rightarrow |A| = 32767$$

$$\frac{(d^2)-1}{4} = \frac{32513-1}{4} = \underbrace{8128}_{\text{Perfect number}} \Rightarrow \underbrace{8191}_{\text{Mersenne prime}} - \underbrace{8128}_{\text{Perfect number}} = \underbrace{63}_{n-1}$$

$$A = \begin{vmatrix} 64 & 1 \\ 65 & 129 \end{vmatrix} \Rightarrow |A| = \underbrace{8191}_{\text{Mersenne prime}}$$

There exists a recurrent triad of functions correlated between them such that their differences generate the Fermat Numbers, this functions are:

$y = 2^x - 1$ (Mersenne Numbers and (Fermat Numbers minus 2), see page 16 (Fermat Numbers with two prime factors)).

$y = 4((2^x - 1)(2^{x-1})) + 1 = 2^{x+1}(2^x - 1) + 1 = d^2$ (squared distance between vectors,).

$y = (2^{x-1})(2^x - 1)$ (contains the Perfect Numbers).

$y = 2^x - 1$	$y = 2^{x+1}(2^x - 1) + 1 = d^2$	$y = (2^{x-1})(2^x - 1)$
1	5	1
3_{F_k-2}	25	6 <i>Perfect number</i>
7	113	28 <i>Perfect number</i>
15_{F_k-2}	481	120
$31_{\text{Mersenne prime}}$	1985	496 <i>Perfect number</i>
...	...	...
9223372036854775807	9223372032559808513	9223372034707292160
($x = 63$)	($x = 31$)	($x = 32$)

In the case of the example above (Mersenne Prime and Perfect Number that looks the same), we have:

$$\begin{array}{c}
 \underbrace{2305843009213693951}_{\text{Mersenne prime}} \\
 \qquad \qquad \qquad \rangle \quad 2147483646 \\
 \qquad \qquad \qquad \qquad \qquad \qquad \rangle \quad \underbrace{1073741823}_{n-1} \\
 \underbrace{2305843007066210305}_{d^2} \\
 \qquad \qquad \qquad \rangle \quad \underbrace{1073741823}_{n-1} \\
 \underbrace{2305843008139952128}_{\text{Perfect number}}
 \end{array}$$


For the matrix of the first non prime Fermat number, we have:

$$A = \begin{vmatrix} 2147483648 & 1 \\ 2147483649 & 4294967297 \end{vmatrix} \Rightarrow |A| = 9223372036854775807$$

$$|A| = 9223372036854775807$$

$$d^2 = 9223372032559808513$$

$$(2^{x-1})(2^x - 1) = 9223372034707292160$$



$7 - 5 = 2_{F_k-3}$. Remember that $2|A|_k + 3 = F_{k+1} \Rightarrow |A|_k = \frac{F_{k+1}-3}{2}$.

Primitive Pythagorean Triples related to Lenghts.

Given the matrix

$$\begin{vmatrix} n & 1 \\ n+1 & 2n+1 \end{vmatrix} \text{ the distance } d = \sqrt{(1-n)^2 + ((2n+1)-(n+1))^2} = \sqrt{2n^2 - 2n + 1}$$

$$\underbrace{(2n-1)^2}_a + \underbrace{(2n^2-2n)^2}_b = \underbrace{(2n^2-2n+1)^2}_c$$

$$|A| - d^2 + 3 = F_k = (2n^2 - 1) - (2n^2 + 2n + 1) + 3 = 2n + 1$$

$$|A| + d^2 + F_k = F_{k+1} = (2n^2 - 1) + (2n^2 - 2n + 1) + (2n + 1) = 4n^2 + 1$$

$$\text{For areas: } b + d^2 = F_{k+1} = (2n^2 + 2n) + (2n^2 - 2n + 1) = 4n^2 + 1$$

This is:

$$|A| + d^2 + F_k = F_{k+1} \Rightarrow |A| + d^2 + |A| - d^2 + 3 = F_{k+1}$$

Comparing polynomials of areas and lenghts we see that n_k related to areas is equal to n_{k+1} related to lenghts.

$$\begin{array}{l} \nearrow \quad 2n^2 - 2n + 1 = d^2 \\ n \\ \searrow \quad 2n^2 + 2n + 1 = \|a\|^2 \end{array}$$

from: $|A|_k = d^2 + F_k - 3$ and knowing that $\mathbf{n}_k$ related to areas is equal to $\mathbf{n}_{k+1}$ related to lenghts, we change values of the lenghts, so we have

$$(2n^2 + 2n + 1)(4n^2 + 4n + 2) = (|\vec{a}|_k |\vec{b}|_k)^2$$

$$(2n^2 - 2n + 1)(4n^2 + 4n + 2) = |A|_{k+1} + 3$$

$$(2n^2 - 2n + 1)(4n^2 - 4n + 2) = 2(d_k)^4$$

Line equations between the two points $(n, n+1)$ and $(1, 2n+1)$.

$$A = \begin{vmatrix} 1 & 1 \\ 2 & 3 \end{vmatrix} \rightarrow |A| = 1$$

$$A = \begin{vmatrix} 2 & 1 \\ 3 & 5 \end{vmatrix} \rightarrow |A| = 7, y = \frac{7}{1} - \frac{2x}{1}$$

$$A = \begin{vmatrix} 8 & 1 \\ 9 & 17 \end{vmatrix} \rightarrow |A| = 127, y = \frac{127}{7} - \frac{8x}{7}$$

$$A = \begin{vmatrix} 128 & 1 \\ 129 & 257 \end{vmatrix} \rightarrow |A| = 32767, y = \frac{32767}{127} - \frac{128x}{127}$$

$$A = \begin{vmatrix} 32768 & 1 \\ 32769 & 65537 \end{vmatrix} \rightarrow |A| = 2147483647, y = \frac{2147483647}{32767} - \frac{32768x}{32767},$$

so, the line equation is:

$$y_k = \frac{|A|_k}{|A|_{k-1}} - \frac{n_k x}{|A|_{k-1}} = \frac{|A|_k}{|A|_{k-1}} - \frac{(|A|_{k-1} + 1)x}{|A|_{k-1}}$$

since all matrices are connected, we can generalize some properties related to (n) for example:

$$1^2 + 1^2 = 2$$

$$\begin{aligned}
2^2 + 2^2 &= 8 \\
8^2 + 8^2 &= 128 \\
128^2 + 128^2 &= 32768 \\
(n_k)^2 + (n_k)^2 &= n_{k+1}
\end{aligned} \tag{3}$$

notice that (n) is the number of *primitive roots* of F_k .

also note that:

$$1 = 2^0$$

$$2 = 2^1$$

$$8 = 2^3$$

$$128 = 2^7$$

$$32768 = 2^{15}$$

0, 1, 3, 7, 15 are the Mersenne Numbers and $|A|$ is a Mersenne Number, this is, the values of $|A|$ are Mersenne Numbers of the form $2n^2 - 1$.

So we can state the next **theorem (2)** for Mersenne Numbers:

If a Mersenne number (Mn_1) have the form $2n^2 - 1$ this is, the equation $(2n^2 - 1 = Mn_1)$ have integer solutions for (n) , the next Mersenne Number Mn_2 doesn't have the form $2n^2 - 1$ and $(2 \cdot Mn_1 + 1 = Mn_2)$ then the value of (n) for $(2n^2 - 1 = Mn_2)$ is equal to $n_1\sqrt{2}$.

$$M_k = 2n^2 - 1; 2M_k + 1 = M_{k+1}$$

$$M_k = 2(n_1)^2 - 1 | n_1 \in Z \Rightarrow \{M_{k+1} = 2(n_2)^2 - 1 | n_2 = n_1\sqrt{2}\}$$

So, for every Fermat Number F_k corresponds a Mersenne Number Mn_k of the form $2n^2 - 1$,


We can see the next table:

$M_{n1} = 2n^2 - 1$	$M_{n2} \neq 2n^2 - 1$
$1 \rightarrow n = 1 = 2^0$	$3 \rightarrow n = 1\sqrt{2}$
$7 \rightarrow n = 2 = 2^1$	$15 \rightarrow n = 2\sqrt{2}$
$31 \rightarrow n = 4 = 2^2$	$63 \rightarrow n = 4\sqrt{2}$
$M_{n1} = 2(2^k)^2 - 1$	$M_{n2} = 2(2^k\sqrt{2})^2 - 1$

Table 2

Angle between vectors.

Example for $F_k = 5$



$$\underbrace{\begin{bmatrix} 2 \\ 3 \end{bmatrix}}_{\vec{a}} \underbrace{\begin{bmatrix} 1 \\ 5 \end{bmatrix}}_{\vec{b}} \rightarrow \begin{aligned} |\vec{a} \cdot \vec{b}| &= 17 \\ \|\vec{a}\| &= \sqrt{13} \\ \|\vec{b}\| &= \sqrt{26} \\ \cos \theta &= \frac{17}{26} \sqrt{2} \end{aligned}$$

$$A = \underbrace{\begin{bmatrix} x_k & 1 \\ y_k & z_k \end{bmatrix}}_{[a_k, b_k, c_k]} \rightarrow \underbrace{\begin{bmatrix} x_{k+1} & 1 \\ y_{k+1} & z_{k+1} \end{bmatrix}}_{[a_{k+1}, b_{k+1}, c_{k+1}]} \quad \begin{aligned} a &= (z)(w) \\ b &= (2)(x)(y) \\ c &= (z)(x) + (y)(w) \end{aligned} \rightarrow b_k - (F_k - 1) = x_{k+1}$$

To obtain n_{k+1} given n_k we have: $(2 \cdot x_k \cdot y_k) - z_k = x_{k+1}$ this is $2 \cdot n \cdot (n+1) - F_k - 1 = n_{k+1} \rightarrow 2(n_k)^2 = n_{k+1}$

To get the matrices and PPT recursively, we have:

$$\begin{aligned} n_k &= 1 \rightarrow 2(n_k)^2 = 2 = n_{k+1} \\ n_{k+1} + 1 &= 3 \\ 2n_{k+1} + 1 &= 5 = a_{k+1} \\ A_k &= \underbrace{\begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}}_{[3,4,5]} b_{k+1} = 2 \cdot n_{k+1} \cdot (n_{k+1} + 1) = 2 \cdot 2 \cdot 3 = 12 \\ c_{k+1} &= b_{k+1} + 1 = 12 + 1 = 13 \\ A_{k+1} &= \underbrace{\begin{bmatrix} 2 & 1 \\ 3 & 5 \end{bmatrix}}_{[5,12,13]} \end{aligned}$$

From the theorem (2) above we can factorize the determinant $|A|$ and obtain it recursively:

$$|A|_{k+1} = [(2|A|_k + 1) - |A|_k \sqrt{2}][(2|A|_k + 1) + |A|_k \sqrt{2}]$$

as for Mersenne Numbers $2Mn_k = Mn_{k+1}$ we have

$$|A|_k = (Mn_k - Mn_{k-1} \sqrt{2})(Mn_k + Mn_{k-1} \sqrt{2})$$

This takes us to:

$$\left\{ \begin{aligned} (F_{k-1} - 1)\sqrt{n+1} &\approx \|a\| \\ (F_{k-1} - 1)\sqrt{F_k + 1} &\approx \|b\| \end{aligned} \right\} \quad \text{Table 3}$$

F_K	$\ a\ $	$(F_{k-1} - 1)\sqrt{\frac{F_k+1}{2}}$	$\ b\ $	$(F_{k-1} - 1)\sqrt{F_k + 1}$
5	3.605551275464	3.46410161513775	5.0990195135928	7.3484692283496
17	12.041594578792	12	17.029386365926	21.213203435597
257	181.72781845386	181.72506706560	257.00194551793	273.06043287155
65537	46341.657113229	46341.657113229	65537.000007629	65793.003898591
4294967297	65536 $\sqrt{2147483649}$	65536 $\sqrt{2147483649}$	65536 $\sqrt{4294967298}$	65536 $\sqrt{4294967298}$

(From Figure 1) Note that triangle (C) have legs $(2n+1 = F_k$ and 1) so hypotenuse $\|b\| \approx F_k$.

Cos θ .

Since $\|b\| = \|a\|\sqrt{2}$, $|a \cdot b| = 2n^2 + 4n + 1$, $\|a\|^2 = 2n^2 + 2n + 1$, $\|b\|^2 = 4n^2 + 4n + 2$ we have:

$$\frac{|a \cdot b|}{\|a\|\|b\|} = \frac{|a \cdot b|}{\|a\|\|a\|\sqrt{2}} = \frac{|a \cdot b|}{\|a\|^2\sqrt{2}} = \frac{2n^2 + 4n + 1}{(2n^2 + 2n + 1)\sqrt{2}}$$

or

$$\frac{|a \cdot b|}{\|a\|\|b\|} = \frac{|a \cdot b|}{\|b\|^2} \sqrt{2} = \frac{(2n^2 + 4n + 1)\sqrt{2}}{4n^2 + 4n + 2}$$

then

$$\cos \theta \approx \lim_{n \rightarrow \infty} \frac{(2n^2 + 4n + 1)\sqrt{2}}{4n^2 + 4n + 2} \approx \lim_{n \rightarrow \infty} \frac{2n^2 + 4n + 1}{(2n^2 + 2n + 1)\sqrt{2}} \approx \frac{1}{\sqrt{2}} \Rightarrow \theta \approx \frac{\pi}{4} \approx 45^\circ$$

note: (the roots of the parabola $(2n^2 - 1)$ are: $-\frac{1}{\sqrt{2}}$ and $\frac{1}{\sqrt{2}}$)

$$|a \cdot b| = \|a\|\|b\| \cos \theta \Rightarrow \cos \theta \|a\|\|b\| - \frac{1}{\sqrt{2}} \|a\|\|b\| = 2n.$$

A PPT conformed by norm values is:

$$a = \sqrt{2\|a\|^2 - 1} = \sqrt{\|b\|^2 - 1} = F_k, b = \|a\|^2 - 1, c = \|a\|^2$$

Primitive Roots, Fermat Primes and their relation with Mersenne Primes and Perfect Numbers.


A Primitive Root modulo (p) is an integer (g) such that $g, g^2, \dots, g^{p-1}$ constitute a reduced residue system modulo (p) , such system is a cyclic group.[4]

From the PPT polynomials, we can relate the *inradius* of a given pythagorean triangle and it's *primitive roots*.

$$\text{number of primitive roots} = \text{inradius} = \frac{ab}{a+b+c} = \frac{(2n+1)(2n^2+2n)}{(2n+1)+(2n^2+2n)+(2n^2+2n+1)} = n$$


$$\text{circumradius} = \frac{c}{2} = n^2 + n + \frac{1}{2}$$

A geometric interpretation is:



$$\Rightarrow \begin{vmatrix} n & 1 \\ n+1 & F_k \end{vmatrix} = Mn_k$$

$\Downarrow$



n is the radius of the circle that fits on the parabola $2n^2 - 1$ at a distance $(0, Mn_k)$ from the origin [5].

We can see the Primitive Roots of the Fermat Primes in the next table. Table 4

F_k	$n = \phi(F_{k_n})$	$\sum \phi(F_{k_n})$	<i>Factorization</i>	<i>Sum of squares</i>	$\sum \phi(F_{k_n}) - \phi(F_{k_n})$
3	1	2	$1 \cdot 2$	$1^2 + 1^2$	2^0
5	2	5	$1 \cdot 5$	$1^2 + 2^2$	$2^0 \cdot 3$
17	8	68	$4 \cdot 17$	$2^2 + 8^2$	$2^2 \cdot 3 \cdot 5$
257	128	16448	$64 \cdot 257$	$8^2 + 128^2$	$2^6 \cdot 3 \cdot 5 \cdot 17$
65537	32768	1073758208	$16384 \cdot 65537$	$128^2 + 32768^2$	$2^{14} \cdot 3 \cdot 5 \cdot 17 \cdot 257$
			$(\phi(F_{k-1_n}))^2 \cdot (F_{k_n})$	$(\phi(F_{k-1_n}))^2 + (\phi(F_{k_n}))^2$	$(\phi(F_{k-1_n}))^2 \cdot \prod_{k=1}^4 F_k$

from (1) page 1, and (3) page 4, we have:

$$\sum \phi(F_{k_n}) = (|A|_k + 1)^2 + (|A|_{k-1} + 1)^2 = \left(\frac{F_k - 1}{2}\right)^2 + \left(\frac{F_{k-1} - 1}{2}\right)^2 = 2^{2^n - 2} \cdot F_k$$

Since the Fermat Primes are connected in relation with their Primitive Roots and the next Fermat Number (4294967297) is not prime and it's number of Primitive Roots is zero, there can't exist more Fermat Primes beyond 65537.

Given the **theorem (2)** "for every Fermat Number F_k corresponds a Mersenne Number Mn_k of the form $2n^2 - 1$ ", we have the next table:

<i>Matrix</i>	<i>Mn</i>	<i>(Mn) Factorization</i>
$\begin{bmatrix} n & 1 \\ n+1 & 2n+1 \end{bmatrix}$	<i>(Determinant)</i>	$[(2n-1) - (n-1)\sqrt{(2)}][(2n-1) + (n-1)\sqrt{(2)}]$
$\begin{bmatrix} \sqrt{2} & 1 \\ 2 + \sqrt{2} & 1 + 2\sqrt{2} \end{bmatrix}$	3	$((2\sqrt{2}-1) - \sqrt{2}(\sqrt{2}-1))((2\sqrt{2}-1) + \sqrt{2}(\sqrt{2}-1))$
$\begin{bmatrix} 2 & 1 \\ 3_{Fp} & 5 \end{bmatrix}$	7	$(3 - 1\sqrt{2})(3 + 1\sqrt{2})$
$\begin{bmatrix} 4 & 1 \\ 5_{Fp} & 9 \end{bmatrix}$	31	$(7 - 3\sqrt{2})(7 + 3\sqrt{2})$
$\begin{bmatrix} 16 & 1 \\ 17_{Fp} & 33 \end{bmatrix}$	511	$(31 - 15\sqrt{2})(31 + 15\sqrt{2})$
$\begin{bmatrix} 256 & 1 \\ 257_{Fp} & 513 \end{bmatrix}$	131071	$(511 - 255\sqrt{2})(511 + 255\sqrt{2})$
$\begin{bmatrix} 65536 & 1 \\ 65537_{Fp} & 131073 \end{bmatrix}$	8589934591	$(131071 - 65535\sqrt{2})(131071 + 65535\sqrt{2})$

$Mn_k(n^2)$ is the Perfect Number related to the given Mersenne Prime (Mn_k).

$Mn_k - (n-1) = Mn_{k-1}$ triangular number.

$Mn_k = (Mn_{k-1} - (n-1)\sqrt{2})(Mn_{k-1} + (n-1)\sqrt{2})$

$Mn_{k+1} = n^2(Mn_k + 1) - 1$.

n is two times the primitive roots of Fermat primes, so, from (1) page 1 and PPT $(F_k, 2n^2 + 2n, 2n^2 + 2n + 1)$, page 3, we have:

$$\left\{ \begin{array}{l} n + 2 \sum \phi(F_{k_n}) = b \\ \Downarrow \\ b - F_k = n_{k+1} - 1 \end{array} \right\} \begin{array}{l} \text{We relate PPT table (page 3)} \\ \text{with Primitive Roots.} \\ \Downarrow \end{array}$$

$2 \cdot b_k - |A|_k = F_{k+1}$, example: PPT $[17, 144, 145] \Rightarrow (2 \cdot 144) - 31 = 257$

We can represent the table this way:

given the values of the function $y = 2^x - 1$

$$\begin{array}{cccccccccc} 2^2 - 1 & 2^3 - 1 & 2^4 - 1 & 2^5 - 1 & 2^6 - 1 & 2^7 - 1 & 2^8 - 1 & 2^9 - 1 & 2^{10} - 1 \\ 3 & 7 & 15 & 31 & 63 & 127 & 255 & 511 & 1023 \\ \checkmark & \checkmark & & \checkmark & & & & \checkmark & \end{array}$$

$$2^x - 1 = |A|_k \Rightarrow 2^{2x-1} - 1 = |A|_{2k}.$$

$$\text{Example: } 2^5 - 1 = 31 \rightarrow 2^{(2 \cdot 5)-1} - 1 = 511$$

The Mersenne Primes and the Perfect Numbers depend on the existence of the Fermat Primes and their primitive roots, every Fermat Prime generates a different number of Mersenne Primes depending on the number of it's Primitive Roots so, there are not infinitely many Mersenne Primes and Perfect Numbers.

The matrices of the table above generate the Mersenne Primes with the primitive roots (2,8,128,32768) of the Fermat Primes and their factors.

$$\begin{aligned}
\begin{bmatrix} 256 & 1 \\ 257_{Fp} & 513 \end{bmatrix} &= {}^{2^{17}-1}_{131071_{Mp}} \\
&\uparrow n^2 \\
\begin{bmatrix} 16 & 1 \\ 17_{Fp} & 33 \end{bmatrix} &= {}^{2^9-1}_{511} \\
&\uparrow n^2 \\
\begin{bmatrix} 4 & 1 \\ 5_{Fp} & 9 \end{bmatrix} &= {}^{2^5-1}_{31}_{Mp} \xrightarrow{n^3} \begin{bmatrix} 2^6 & 1 \\ 64 & 129 \end{bmatrix} = {}^{2^{13}-1}_{8191_{Mp}} \\
&\uparrow n^2 \qquad \qquad \qquad \uparrow n^2 \\
\begin{bmatrix} 2 & 1 \\ 3_{Fp} & 5_{Fp} \end{bmatrix} &= {}^{2^3-1}_7_{Mp} \xrightarrow{n^3} \begin{bmatrix} 8 & 1 \\ 9 & 17_{Fp} \end{bmatrix} = {}^{2^7-1}_{127}_{Mp} \xrightarrow{n^3} \begin{bmatrix} 2^9 & 1 \\ 512 & 1025 \end{bmatrix} = {}^{2^{19}-1}_{524287_{Mp}} \\
&\uparrow n^2 \qquad \qquad \qquad \downarrow 2n^2 \\
\begin{bmatrix} \sqrt{2} & 1 \\ 1+\sqrt{2} & 1+2\sqrt{2} \end{bmatrix} &= 3 \begin{bmatrix} 128 & 1 \\ 129 & 257_{Fp} \end{bmatrix} = {}^{2^{15}-1}_{32767} \\
&\qquad \qquad \qquad \downarrow 2n^2 \\
\begin{bmatrix} 2^{15} & 1 \\ 32768 & 65537_{Fp} \end{bmatrix} &= 2147483647_{Mp} \xrightarrow{8n^4} \begin{bmatrix} 2^{63}=2 \cdot 2147483648^2 & 1 \\ 9223372036854775808 & 18446744073709551617 \end{bmatrix} \\
&\qquad \qquad \qquad \downarrow n^2 \\
\begin{bmatrix} 2^{30} & 1 \\ 1073741824 & 2147483649 \end{bmatrix} &= {}^{2^{61}-1}_{2305843009213693951_{Mp}} \\
&\downarrow 128^2 n = 16384 n = 2^{14} n \\
\begin{bmatrix} 2^{44} & 1 \\ 17592186044416 & 35184372088833 \end{bmatrix} &= {}^{2^{89}-1}_{618970019642690137449562111_{Mp}} \\
&\downarrow 8^3 n = 512 n = 2^9 n \\
\begin{bmatrix} 2^{53} & 1 \\ 9007199254740992 & 18014398509481985 \end{bmatrix} &= {}^{2^{107}-1}_{162259276829213363391578010288127_{Mp}}
\end{aligned}$$

Note that when n is a perfect square, the Mersenne Prime finishes in 1, otherwise in 7.

We can find other relations between Mersenne Primes.

$$\begin{aligned} & \left(\frac{\binom{2^{44}}{17592186044416}}{\binom{M_p+1=2^{13}}{8192}} \right) - 1 = 2147483647_{M_p} \\ & \quad \Downarrow \\ & \binom{M_p+1=2^{31}}{2147483648} \binom{2^{30}}{1073741824} - 1 = 2305843009213693951_{M_p} = \binom{M_p+1=2^{17}}{131072} \binom{2^{44}}{17592186044416} - 1 \\ & \left(\frac{\binom{M_p+1=2^{107}}{162259276829213363391578010288128}}{\binom{M_p+1=2^{61}}{2305843009213693952}} \right) = \left(\frac{\binom{M_p+1=2^{61}}{2305843009213693952}}{\binom{2^{15}}{32768}} \right) \end{aligned}$$

To connect this table with the *primitive roots* we check the *primitive roots tables* of the Fermat Primes.

$F_k = 3$ Root{2}	$\sum \bmod 3$	$F_k = 5$ Roots{2,3}		$\sum \bmod 5$
(2)		(2)	(3)	
1	1	1	1	2
2	2 (Prim root.)	2	3	5 (Primitive roots.)
$\sum = 3$		4	4	8
		3	2	5 (Primitive roots.)
		$\sum = 10$	$\sum = 10$	

$F_k = 17$ Primitive Roots {3,5,6,7,10,11,12,14}

(3)	(5)	(6)	(7)	(10)	(11)	(12)	(14)	$\sum \bmod 17$
1	1	1	1	1	1	1	1	8
3	5	6	7	10	11	12	14	68 (Primitive roots.)
9	8	2	15	15	2	8	9	68
10	6	12	3	14	5	11	7	68 (Primitive roots.)
13	13	4	4	4	4	13	13	68
5	14	7	11	6	10	3	12	68 (Primitive roots.)
15	2	8	9	9	8	2	15	68
11	10	14	12	5	3	7	6	68 (Primitive roots.)
16	16	16	16	16	16	16	16	128
14	12	11	10	7	6	5	3	68 (Primitive roots.)
8	9	15	2	2	15	9	8	68
7	11	5	14	3	12	6	10	68 (Primitive roots.)
4	4	13	13	13	13	4	4	68
12	3	10	6	11	7	14	5	68 (Primitive roots.)
2	15	9	8	8	9	15	2	68
6	7	3	5	12	14	10	11	68 (Primitive roots.)

$\sum = F_k \left(\frac{F_k-1}{2} \right) = 136$ for all roots.

Note that $\{3^n, 5^n, 6^n, 7^n, 10^n, 11^n, 12^n, 14^n\} \bmod(17)$ are cyclic, so the entire system is cyclic, the summation of all the residues are $\{3, 10, 136, 32896, 2147516416\}$

and from **(3) pages (4,5)** they are connected this way:

$$3^2 + 1^2 = 3^2 + (2-1)^2 = 2^1 \cdot 5 = 10$$

$$10^2 + 6^2 = 10^2 + (8-2)^2 = 2^3 \cdot 17 = 136$$

$$136^2 + 120^2 = 136^2 + (128-8)^2 = 2^7 \cdot 257 = 32896$$

$$32896^2 + 32640^2 = 32896^2 + (32768-128)^2 = 2^{15} \cdot 65537 = 2147516416$$

(2, 8, 128, 32768) is the summation of the horizontal line with equal values

(excepting the first line) on every *Primitive root* and is $\frac{F_{(k+1)}-1}{2}$.

since $2|A| + 3 = F_{(k+1)}$ we have:

$$\frac{10}{3^2} = \frac{7+3}{3^2} \Rightarrow \frac{136}{10^2} = \frac{31+3}{5^2} \Rightarrow \frac{32896}{136^2} = \frac{511+3}{17^2} \Rightarrow \frac{2147516416}{32896^2} = \frac{131071+3}{257^2}$$

(7,31,511,131071) are the values of the determinants of the matrices that start the table.

Sigma(n)

Knowing that $n = \phi(Fk_n)$ we have:

$$\sigma(1) = 1 \Rightarrow \sigma(1) + 2 = 3$$

$$\sigma(2) = 3 \Rightarrow \sigma(2) + 2 = 5$$

$$\sigma(8) = 15 \Rightarrow \sigma(8) + 2 = 17$$

$$\sigma(128) = 255 \Rightarrow \sigma(128) + 2 = 257$$

$$\sigma(32768) = 65535 \Rightarrow \sigma(32768) + 2 = 65537$$

$$\text{Since } \left\{ \prod_1^k F_k \right\} + 2 = F_{k_n} \Rightarrow \sigma(n) = \left\{ \prod_1^k F_k \right\} \Rightarrow \sigma\left(\frac{F_k-1}{2}\right) = F_k - 2$$

$$\text{thus } \sigma(\phi(F_{k_n})) = F_k - 2$$

$$\text{Given } (n) = \{1, 2, 8, 128, 32768\} \Rightarrow (2n_k)^2 = 2n_{(k+1)} \Rightarrow \sigma(2n_k)^2 = \sigma_{(k+1)}$$

Fermat Factors.

By now we will assume that all Fermat Factors have the next form:

$$\begin{vmatrix} n & 1 \\ n+1 & F_k \end{vmatrix} \text{ where } F_k = f_1 \cdot f_2 \cdot f_3 \cdots f_k \text{ then } \forall f_k \{f_k \equiv 1 \pmod{24}\}$$

or $f_k \equiv 17 \pmod{24}$, from sigma properties and for $F_k = f_1 \cdot f_2$ we have:

$$n = \left(\frac{f_2-1}{2}\right)(f_1) + \left(\frac{f_1-1}{2}\right) = \left(\frac{f_1-1}{2}\right)(f_2) + \left(\frac{f_2-1}{2}\right)$$

This takes us to:

$$\sigma(F_k) - F_k - f_1 - 1 = f_2$$

$$\text{Example: } F_k = 4294967297 = (641 \cdot 6700417), \sigma(F_k) = 4301668356$$

$$4301668356 - 4294967297 - 6700417 - 1 = 641$$

so, given any two factors f_1 and f_2 of the F_k sequence, we have:

$$f_1 = (x_1 \cdot 24) + 17 \text{ and } f_2 = (x_2 \cdot 24) + 1, \text{ then:}$$

$$f_1 = (x_1 - f_2 + 24x_2 + 17)24 + 17 \text{ and } f_2 = (x_2 - f_1 + 24x_1 + 1)24 + 1$$

knowing that $\sigma(n) = \prod_{i=1}^k (1 + p_i + p_i^2 + \cdots + p_i^{\alpha_i})$ we have for a two factor F_k

$$\sigma(n) = (24x_1 + 18)(24x_2 + 2) = 576x_1x_2 + 48x_1 + 432x_2 + 36$$

$$\text{example: } \sigma(F_k) = (576 \cdot 26 \cdot 279184) + (48 \cdot 26) + (432 \cdot 279184) + 36 = 4301668356$$

notice that for a Fermat Number with three factors $F_k = f_1 \cdot f_2 \cdot f_3$, two of them will be $f_1 \equiv 17 \pmod{24}$ and the other $f_2 \equiv 1 \pmod{24}$ or viceversa, but them all can't have the same residue. this applies for every F_k .

Fermat Numbers with two Prime Factors.

Given a Fermat Factor $\mathbf{f} \equiv 1 \pmod{4}$ we have $f = x^2 + y^2$, for a

Fermat Number with two factors $\mathbf{f}_1 \cdot \mathbf{f}_2$ then:

$$A = \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} \Rightarrow |\vec{a} \cdot \vec{b}| = F_{k-1} - 1$$

$$\text{if } f^2 = x^2 + y^2$$

$$A = \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} \Rightarrow |\vec{a} \cdot \vec{b}| = F_k - 2$$

$$\text{since } F_k = (F_{k-1} - 1)^2 + 1 \Rightarrow F_k = \left(|\vec{a} \cdot \vec{b}|_{k-1}\right)^2 + 1$$

This rules are:

$$F_k = x^2 + y^2 \text{ such that } x + y = F_{k-1}$$

$$(F_k)^2 = (x')^2 + (y')^2 \text{ such that } x' = 2y \wedge y' = y^2 - 1$$

Example:

$$F_k = 4294967297 = 641 \cdot 6700417$$

$$641 = 4^2 + 25^2 \text{ and } 641^2 = 200^2 + 609^2$$

$$6700417 = 409^2 + 2556^2 \text{ and } 6700417^2 = 2090808^2 + 6365855^2$$

so, we have:

$$A = \begin{bmatrix} 4 & 409 \\ 25 & 2556 \end{bmatrix} \Rightarrow |\vec{a} \cdot \vec{b}| = 65536$$

and

$$A = \begin{bmatrix} 2090808 & 200 \\ 6365855 & 609 \end{bmatrix} \Rightarrow |\vec{a} \cdot \vec{b}| = 4294967295$$

for $f^2 = x^2 + y^2$, since $|\vec{a} \cdot \vec{b}| \cos(\theta) = f_1 \cdot f_2 \cos(\theta) \Rightarrow \cos(\theta) = \frac{F_k - 2}{F_k}$
since we have the recurrence:

$$A_k = \begin{bmatrix} n & 1 \\ n+1 & F_k \end{bmatrix} \Rightarrow A_{k+1} = \begin{bmatrix} |A_k| & 1 \\ |A_k| + 1 & F_{k+1} \end{bmatrix}$$

$$\cos(\theta) = \frac{F_k - 2}{F_k} = \frac{1}{1 + \frac{1}{|A_{k-1}| + \frac{1}{2}}}$$

Fermat Numbers with more than two Prime Factors.

Given a Fermat Number $F_k = f_1 \cdot f_2 \cdots f_k$ such that $\underbrace{\begin{bmatrix} x_1 \\ y_1 \end{bmatrix}}_{f_1}, \underbrace{\begin{bmatrix} x_2 \\ y_2 \end{bmatrix}}_{f_2}, \underbrace{\begin{bmatrix} x_k \\ y_k \end{bmatrix}}_{f_k}$ where

$$(f_1)^2 = (x_1)^2 + (y_1)^2, (f_2)^2 = (x_2)^2 + (y_2)^2, (f_k)^2 = (x_k)^2 + (y_k)^2$$

we have

$$A = \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} \Rightarrow |\vec{a} \cdot \vec{b}|_1 = x_1 \cdot x_2 + y_1 \cdot y_2 = (f_{1.2})^2 \Rightarrow (f_{1.2})^2 = (x_{1.2})^2 + (y_{1.2})^2$$

then

$$A_k = \begin{bmatrix} x_{1.2} & x_k \\ y_{1.2} & y_k \end{bmatrix} \Rightarrow |\vec{a} \cdot \vec{b}|_k \Rightarrow \frac{f_1 \cdot f_2 \cdots f_k}{|\vec{a} \cdot \vec{b}|_k} = \frac{(f_1 \cdot f_2 \cdots f_k) - |\vec{a} \cdot \vec{b}|_k}{|\vec{a} \cdot \vec{b}|_k} + 1$$

The John Wallis Pairs.

The *Wallis Pairs* (W_x, W_y) are numbers of the form $\sigma(x^2) = \sigma(y^2)$, here are some of them representing the 'y' side of a primitive pythagorean triangle where the hypotenuse is a pythagorean prime.

$$[12, 35, 37] \Rightarrow 37 = 1^2 + 6^2; (1 + 6) \cdot 5 = 35$$

$$[28, 45, 53] \Rightarrow 53 = 2^2 + 7^2; (2 + 7) \cdot 5 = 45$$

$$[48, 55, 73] \Rightarrow 73 = 3^2 + 8^2; (3 + 8) \cdot 5 = 55$$

$$[72, 65, 97] \Rightarrow 97 = 4^2 + 9^2; (4 + 9) \cdot 5 = 65$$

$$[132, 85, 157] \Rightarrow 157 = 6^2 + 11^2; (6 + 11) \cdot 5 = 85$$

$$[168, 95, 193] \Rightarrow 193 = 7^2 + 12^2; (7 + 12) \cdot 5 = 95$$

Since every $F_k \equiv 2 \pmod{5}$ and for every Fermat Number with two prime factors:

$$f^2 = x^2 + y^2 \Rightarrow \begin{vmatrix} x_x & y_1 \\ x_2 & y_2 \end{vmatrix} \Rightarrow |\vec{a} \cdot \vec{b}| = F_k - 2 \text{ then } F_k - 2 \text{ is } W_y \text{ part of a}$$

Wallis Pair.

For the *Fermat Primes* we have:

$$\begin{array}{lll}
[4, 3, 5] & 5 = \underbrace{1^2 + 2^2}_{x+y=3} & (1+2) \cdot 1 = 3 \\
[8, 15, 17] & 17 = \underbrace{1^2 + 4^2}_{x+y=5} & (1+4) \cdot 3 = 15 \\
[32, 255, 257] & 257 = \underbrace{1^2 + 16^2}_{x+y=17} & (1+16) \cdot 15 = 255 \\
[512, 65535, 65537] & 65537 = \underbrace{1^2 + 256^2}_{x+y=257} & (1+256) \cdot 255 = 65535
\end{array}$$

So the recurrence of the 'y' part of the *Wallis Pairs* is next:

$$\frac{F_k-2}{5} = \frac{|\vec{a} \cdot \vec{b}|}{5} = \frac{4294967295}{5} = \underbrace{858993459}_{3 \cdot 17 \cdot 257 \cdot 65537} \dots 1$$

$$\frac{858993459}{65537} = \frac{65535}{5} = \underbrace{13107}_{3 \cdot 17 \cdot 257} \dots 1$$

$$\frac{13107}{257} = \frac{255}{5} = \underbrace{51}_{3 \cdot 17} \dots 1$$

$$\frac{51}{17} = \frac{15}{5} = 3 \dots 1$$

What $\dots 1$ means? The W_y part of the *Wallis Pair* is a vector product and is related to the *cosine* of an angle, the missing $\dots 1$ appears as the last digit of a *Mersenne Number* in the table of page 12 as (*Determinant*).

For the W_x part of the *Wallis Pair*:

$$y = 65535, x = 52428 \dots 7 \Rightarrow 65535 - 52428 = 13107$$

$$y = 255, x = \frac{52428}{257} = 204 \dots 7 \Rightarrow 255 - 204 = 51$$

$$y = 15, x = \frac{204}{17} = 12 \dots 7 \Rightarrow 15 - 12 = 3$$

$$\underbrace{\frac{4}{5} \prod_1^n F_k}_{W_x}, \underbrace{\prod_1^n F_k}_{W_y}$$

Again, $\dots 7$ is missing, but this time it is related to the *Mersenne Numbers* whose last digit is 7, this is 127,2047,524287.

The *Wallis Pairs* are generated by a first pair, the $(W_x = 4, W_y = 5)$, in the same way, we can generate *Pythagorean Triples (not primitive)* with the triple $[3, 4, 5]$, so we have:

$$[\underbrace{39321}_a, \underbrace{52428}_b, \underbrace{65535}_c]$$

$$[\underbrace{153}_a, \underbrace{204}_{W_x}, \underbrace{255}_{W_y}]$$

$$[9, 12, 15]$$

$$[3, 4, 5]$$

$$\text{where } c - b = b - a; \frac{a_k}{a_{k-1}} = F_{k-1}$$

since to form a *Fibonacci Box* we have: $\frac{c-a}{b}, \frac{c-b}{a}$ then all these *Pythagorean*

Triples are generated by the same matrix, $A = \begin{vmatrix} 1 & 1 \\ 2 & 3 \end{vmatrix}$ this is the first *Fibonacci Box* with members (1, 1, 2, 3), this *Fibonacci Box* generates the triple (3, 4, 5) and happens that $|A| = 1$ and $|a \cdot b| = 7$.

Cotes Theorem.

Given a unit circle with a regular polygon inscribed inside and a point located at the 'x axis' then, the product of the distances from the point to the polygon vertices is equal to $r^N - x^N$ when the point is inside the circle or $x^N - r^N$ when the point is outside the circle, being $r = 1$ and $N =$ number of polygon vertices.

$$\text{if } x^n = F_k - 1 \Rightarrow x^n - 1 = F_k - 2 = \left\{ \prod_1^k F_k \right\}$$

The point 'x' is located at (2, 0), the distance from point 'x' to the point located at $\angle = 180^\circ = 3$.

To find the factors of F_k we factorize F_{k+1} . So, we have:

3

$$\sqrt{5} \cdot \sqrt{5} = 5 \quad (N = 4)$$

$$(5 - 2\sqrt{2}) \cdot (5 + 2\sqrt{2}) = 17 \quad (N = 8)$$

For $(N = 16)$:

$$(5 - 2\sqrt{2 - \sqrt{2}}) \cdot (5 - 2\sqrt{2 + \sqrt{2}}) \cdot (5 + 2\sqrt{2 - \sqrt{2}}) \cdot (5 + 2\sqrt{2 + \sqrt{2}}) = 257$$

For $(N = 32)$:

$$\angle = 11.25^\circ \Rightarrow d = (5 - 2\sqrt{2 + \sqrt{2 + \sqrt{2}}})$$

$$\angle = 22.5^\circ \Rightarrow d = (5 - 2\sqrt{2 + \sqrt{2}})_{(257)}$$

$$\angle = 33.75^\circ \Rightarrow d = (5 - 2\sqrt{2 + \sqrt{2 - \sqrt{2}}})$$

$$\angle = 45^\circ \Rightarrow d = (5 - 2\sqrt{2})_{(17)}$$

$$\angle = 56.25^\circ \Rightarrow d = (5 - 2\sqrt{2 - \sqrt{2 - \sqrt{2}}})$$

$$\angle = 67.5^\circ \Rightarrow d = (5 - 2\sqrt{2 - \sqrt{2}})_{(257)}$$

$$\angle = 78.75^\circ \Rightarrow d = (5 - 2\sqrt{2 - \sqrt{2 + \sqrt{2}}})$$

$$\angle = 90^\circ \Rightarrow d = (\sqrt{5})_{(5)}$$

$$\angle = 101.25^\circ \Rightarrow d = (5 + 2\sqrt{2 - \sqrt{2 + \sqrt{2}}})$$

$$\angle = 112.5^\circ \Rightarrow d = (5 + 2\sqrt{2 - \sqrt{2}})_{(257)}$$

$$\angle = 123.75^\circ \Rightarrow d = (5 + 2\sqrt{2 - \sqrt{2 - \sqrt{2}}})$$

$$\angle = 135^\circ \Rightarrow d = (5 + 2\sqrt{2})_{(17)}$$

$$\angle = 146.25^\circ \Rightarrow d = \left(5 + 2\sqrt{2 + \sqrt{2 - \sqrt{2}}}\right)$$

$$\angle = 157.5^\circ \Rightarrow d = \left(5 + 2\sqrt{2 + \sqrt{2}}\right)_{(257)}$$

$$\angle = 168.75^\circ \Rightarrow d = \left(5 + 2\sqrt{2 + \sqrt{2 + \sqrt{2}}}\right)$$

$$\angle = 180^\circ \Rightarrow d = 3$$

So:

$$\begin{aligned} & \left(5 - 2\sqrt{2 + \sqrt{2 + \sqrt{2}}}\right) \cdot \left(5 - 2\sqrt{2 + \sqrt{2 - \sqrt{2}}}\right) \cdot \left(5 - 2\sqrt{2 - \sqrt{2 - \sqrt{2}}}\right) \cdot \left(5 - 2\sqrt{2 - \sqrt{2 + \sqrt{2}}}\right) \cdot \\ & \left(5 + 2\sqrt{2 - \sqrt{2 + \sqrt{2}}}\right) \cdot \left(5 + 2\sqrt{2 - \sqrt{2 - \sqrt{2}}}\right) \cdot \left(5 + 2\sqrt{2 + \sqrt{2 - \sqrt{2}}}\right) \cdot \left(5 + 2\sqrt{2 + \sqrt{2 + \sqrt{2}}}\right) \\ & = 65537 \end{aligned}$$

Given this pattern, the next Fermat Number should have this form:

$$\begin{aligned} & \left(5 + 2\sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2}}}}\right) \cdot \left(5 + 2\sqrt{2 - \sqrt{2 + \sqrt{2 + \sqrt{2}}}}\right) \cdot \left(5 + 2\sqrt{2 + \sqrt{2 - \sqrt{2 + \sqrt{2}}}}\right) \cdots \\ & \cdots \left(5 - 2\sqrt{2 - \sqrt{2 - \sqrt{2 - \sqrt{2}}}}\right) = 4294967297 \end{aligned}$$

Knowing that:

$$F_{k-2} = \left\{ \prod_1^k F_k \right\} = x^2 - 1 \prod_{n=1}^{\frac{N}{2}-1} \{1 + x^2 - 2x \cos(\frac{2\pi n}{N})\} = 3 \prod_{n=1}^{\frac{N}{2}-1} \{5 - 4 \cos(\frac{2\pi n}{N})\}$$

$$\Rightarrow 2\sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2 + \cdots}}}} = 4 \cos\left(\frac{2\pi n}{N}\right)$$

This is:

$$\begin{aligned} & \sqrt{5} \cdot \sqrt{5} = (5 - 4 \cos(\frac{2\pi}{4})) = 5 \\ & (5 - 2\sqrt{2}) \cdot (5 + 2\sqrt{2}) = (5 - 4 \cos(\frac{2\pi}{8})) \cdot (5 - 4 \cos(\frac{6\pi}{8})) = 17 \\ & (5 - 2\sqrt{2 + \sqrt{2}}) \cdot (5 - 2\sqrt{2 - \sqrt{2}}) \cdot (5 + 2\sqrt{2 - \sqrt{2}}) \cdot (5 + 2\sqrt{2 + \sqrt{2}}) \\ & = (5 - 4 \cos(\frac{2\pi}{16})) \cdot (5 - 4 \cos(\frac{6\pi}{16})) \cdot (5 - 4 \cos(\frac{10\pi}{16})) \cdot (5 - 4 \cos(\frac{14\pi}{16})) = 257 \\ & (5 - 4 \cos(\frac{2\pi}{32})) \cdot (5 - 4 \cos(\frac{6\pi}{32})) \cdot (5 - 4 \cos(\frac{10\pi}{32})) \cdot (5 - 4 \cos(\frac{14\pi}{32})) \\ & \cdot (5 - 4 \cos(\frac{18\pi}{32})) \cdot (5 - 4 \cos(\frac{22\pi}{32})) \cdot (5 - 4 \cos(\frac{26\pi}{32})) \cdot (5 - 4 \cos(\frac{30\pi}{32})) = 65537 \end{aligned}$$

As corollary:

$$\text{Since Fermat } F_k = 3 \prod_{n=1}^{\frac{N}{2}-1} \{5 - 4 \cos(\frac{2\pi n}{N})\} + 2, \text{ Mersenne } M_k = \prod_{n=1}^{\frac{N-1}{2}} \{5 - 4 \cos(\frac{2\pi n}{N})\}$$

and we know that $2|A|_k + 3 = F_k, \forall |A|_k = M_k$

$$\text{then: } 3 \prod_{n=1}^{\frac{N}{2}-1} \{5 - 4 \cos(\frac{2\pi n}{N})\} + 2 = 2 \prod_{n=1}^{\frac{N-1}{2}} \{5 - 4 \cos(\frac{2\pi n}{N})\} + 3 \quad (N_{F_k} \neq N_{M_k})$$

since $\frac{N-1}{2} - (\frac{N}{2} - 1) = \frac{1}{2}$ this equation is just a *switch* of the cosine values.

Example:

$$3(5 - 4 \cos(\frac{2\pi}{4})) + 2 = 2(5 - 4 \cos(\frac{2\pi}{3})) + 3 = 17$$

Note that for every F_k we have the Fermat Number via Cote's Theorem,

Fermat Number as a product and Mersenne Number as a product, here we have

$F_k = 17$:

$3 \left(5 - 4 \cos \left(\frac{2\pi}{4}\right)\right) + 2 = 17$ *Cote's Theorem.*

$\left(5 - 4 \cos \left(\frac{2\pi}{8}\right)\right) \cdot \left(5 - 4 \cos \left(\frac{6\pi}{8}\right)\right) = 17$ F_k *as product.*

$\left(5 - 4 \cos \left(\frac{2\pi}{3}\right)\right) = 7$ M_k *as product.*

$2|A| + 3$ is equivalent to: $\frac{N}{2} - 1 \Rightarrow \frac{N-1}{2}$

Notice that (from (2) page two), when the polynomial $2n^2 + 4n + 1$ takes the values of $n = |A|_k$ we obtain $|A_{k+1}|$

this is: $2n^2 + 4n + 1 = M_k \Rightarrow 2M_k + 3 = F_k$

Example: $2 \cdot 7^2 + 4 \cdot 7 + 1 = 127 \Rightarrow 2 \cdot 127 + 3 = 257 \Rightarrow 2 \cdot 127^2 + 4 \cdot 127 + 1 = 32767 \Rightarrow 2 \cdot 32767 + 3 = 65537$

This is the system we already obtained:

$$A_1 = \begin{vmatrix} 2 & 1 \\ 3 & 5 \end{vmatrix} \Rightarrow |A_1| = 7 \Rightarrow A_2 = \begin{vmatrix} 8 & 1 \\ 9 & 17 \end{vmatrix} \Rightarrow |A_2| = 127 \Rightarrow A_3 = \begin{vmatrix} 128 & 1 \\ 129 & 257 \end{vmatrix}$$

$\Rightarrow |A_3| = 32767$

then: $2^k - 1 = M_k \Rightarrow 2(2^k - 1) + 3 = 2^{k+1} + 1 = F_k \Rightarrow 2(2^k - 1) + 3$

$= 2^{k+1} - 2 + 3 = 2^{k+1} + 1$

since $M_k = \frac{F_k - 3}{2} \Rightarrow \frac{F_k - 1}{2} = M_k + 1 \Rightarrow 3^{\frac{F_k - 1}{2}} = 3^{M_k + 1}$

Pepin's Test:

$3^{\frac{F_k - 1}{2}} \equiv -1 \pmod{F_k} \Rightarrow 3^{M_k + 1} \equiv -1 \pmod{2M_k + 3}$

Pepin's Test as Cote's Theorem:

$$8 \prod_{n=1}^N \left\{ 10 - 6 \cos \left(\frac{2\pi n}{\left(\frac{F_k - 1}{2}\right)} \right) \right\} \equiv -1 \pmod{\left(3 \prod_{n=1}^{\frac{N}{2} - 1} \left\{ 5 - 4 \cos \left(\frac{2\pi n}{N} \right) \right\} + 2 \right)}$$

$$\text{since: } 3 \prod_{n=1}^{\frac{N}{2} - 1} \left\{ 5 - 4 \cos \left(\frac{2\pi n}{N} \right) \right\} + 2 = 2 \prod_{n=1}^{\frac{N-1}{2}} \left\{ 5 - 4 \cos \left(\frac{2\pi n}{N} \right) \right\} + 3$$

$$\text{then: } \left[\frac{8 \prod_{n=1}^N \left\{ 10 - 6 \cos \left(\frac{2\pi n}{\left(\frac{F_k - 1}{2}\right)} \right) \right\} + 2}{2 \prod_{n=1}^{\frac{N-1}{2}} \left\{ 5 - 4 \cos \left(\frac{2\pi n}{N} \right) \right\} + 3} \right] = \left[\frac{8 \prod_{n=1}^N \left\{ 10 - 6 \cos \left(\frac{2\pi n}{\left(\frac{F_k - 1}{2}\right)} \right) \right\} + 2}{3 \prod_{n=1}^{\frac{N}{2} - 1} \left\{ 5 - 4 \cos \left(\frac{2\pi n}{N} \right) \right\} + 2} \right] \in \mathbb{Z} \quad \{\forall F_k \text{ prime}\}$$

For $F_k = 17$ we have:

$$\left[\frac{((10 - 3\sqrt{2}) \cdot (10 + 3\sqrt{2}) \cdot 10 \cdot 4 \cdot 2) + 2}{(5 - 2\sqrt{2}) \cdot (5 + 2\sqrt{2})} \right] = \frac{8 \cdot (10 - 6 \cos \left(\frac{2\pi}{8}\right)) \cdot (10 - 6 \cos \left(\frac{4\pi}{8}\right)) \cdot (10 - 6 \cos \left(\frac{6\pi}{8}\right)) + 2}{3 \left((5 - 4 \cos \left(\frac{2\pi}{4}\right)) + 2 \right)}$$

$$= \frac{8 \left((10 - 6 \cos \left(\frac{2\pi}{8}\right)) \cdot (10 - 6 \cos \left(\frac{4\pi}{8}\right)) \cdot (10 - 6 \cos \left(\frac{6\pi}{8}\right)) \right) + 2}{(5 - 4 \cos \left(\frac{2\pi}{8}\right)) \cdot (5 - 4 \cos \left(\frac{6\pi}{8}\right))} \in \mathbb{Z}$$

this is:

$$\frac{386 \times \left((5 - 4 \cos \left(\frac{2\pi}{8}\right)) \cdot (5 - 4 \cos \left(\frac{6\pi}{8}\right)) \right)}{(5 - 4 \cos \left(\frac{2\pi}{8}\right)) \cdot (5 - 4 \cos \left(\frac{6\pi}{8}\right))}$$

this condition is not infinite and have a breakpoint.

Pepin's test in Mersenne numbers terms:

$$2 \prod_{n=1}^{M_k + 1} \left\{ 10 - 6 \cos \left(\frac{2\pi n}{2M_k + 3} \right) \right\} \equiv -1 \pmod{\left(2 \prod_{n=1}^{\frac{N-1}{2}} \left\{ 5 - 4 \cos \left(\frac{2\pi n}{N} \right) \right\} + 3 \right)}$$

Multiplying different values of (n).

Given consecutive Mersenne Primes and their respective matrices:

$$\left| \begin{array}{cc} n_1 & 1 \\ n_1 + 1 & 2n_1 + 1 \end{array} \right|, \left| \begin{array}{cc} n_2 & 1 \\ n_2 + 1 & 2n_2 + 1 \end{array} \right|, \dots, \left| \begin{array}{cc} n_k & 1 \\ n_k + 1 & 2n_k + 1 \end{array} \right|$$

$|A_1|=2^{p_1}-1=7 \quad |A_2|=2^{p_2}-1=31 \quad |A_k|=2^{p_k}-1=Mp_k$

Then:

$$\left(\frac{n_2}{n_1} \right) \cdot \left(\frac{n_3}{n_2} \right) \dots \left(\frac{n_k}{n_{k-1}} \right) = 2^{\frac{p_k-3}{2}}$$

$$\text{This is } \frac{F_k-3}{2} = Mp_k \Rightarrow \frac{2^{2^{n-1}}+1-3}{2} = 2^{2^{n-1}-1} - 1$$

Example:

Given the next matrices and their determinants.

$$\left| \begin{array}{cc} 2 & 1 \\ 3 & 5 \end{array} \right|; \left| \begin{array}{cc} 4 & 1 \\ 5 & 9 \end{array} \right|; \left| \begin{array}{cc} 8 & 1 \\ 9 & 17 \end{array} \right|; \left| \begin{array}{cc} 64 & 1 \\ 65 & 129 \end{array} \right| \Rightarrow \left(\frac{4}{2} \right) \cdot \left(\frac{8}{4} \right) \cdot \left(\frac{64}{8} \right) = 2 \cdot 2 \cdot 8 = 2^{\frac{13-3}{2}} = 2^5 = 32$$

$|A|=7 \quad |A|=31 \quad |A|=127 \quad |A|=8191$

The number 13 is ' p ' of $2^p - 1 = \frac{2^{13}-1}{8191}$

Note that from *Pepin's Test*: $3^{\frac{F_k-1}{2}} \equiv -1 \pmod{F_k}$

$$\text{from } \left| \begin{array}{cc} n & 1 \\ n+1 & F_k \end{array} \right| \Rightarrow \frac{F_k-1}{2} = n, \text{ Pepin's Test turns to } 3^n \equiv -1 \pmod{F_k}$$

this turns to $F_k \equiv 3 \pmod{n-1}$ where $\frac{F_k-3}{n-1} = 2$

then $F_k \equiv 3 \pmod{(|A|_{k-1})} \Rightarrow F_{(k+1)} = 2|A| + 3$

32768

$$\text{Given a matrix } A = \left| \begin{array}{cc} n & 1 \\ n+1 & 2n+1 \end{array} \right| \Rightarrow \left| \begin{array}{cc} n & 1 \\ n+1 & F_k \end{array} \right| \Rightarrow \left| \begin{array}{cc} 32768 & 1 \\ 32769 & 65537 \end{array} \right|$$

$|A| = \frac{2^{31}-1}{2147483647} M_p$ being **n=32768** the inradius of the Pythagorean Triangle with sides $\underbrace{65537}_{2n+1}, \underbrace{2147549184}_{2n^2+2n}, \underbrace{2147549185}_{2n^2+2n+1}$. The inradius 32768 is connected with

all the Mersenne Primes as we can see next.

$Mp_k < 32768$

$$32768 = [(4681)(7)] + 1$$

$$32768 = [(1057)(31)] + 1$$

$$32768 = [(258)(127)] + 2$$

$$32768 = [(4)(8191)] + 4$$

$$Mp_k > 32768 \quad [(4^n - 1)(32768)] + (32768 - 1)$$

$$M_p = \frac{2^{17}-1}{131071} = \left[\left(\frac{2^2-1}{3} \right) (32768) \right] + (32768 - 1)$$

$$M_p = \frac{2^{19}-1}{524287} = \left[\left(\frac{2^4-1}{15} \right) (32768) \right] + (32768 - 1)$$

$$M_p = \frac{2^{31}-1}{2147483647} = \left[\left(\frac{2^{16}-1}{65535} \right) (32768) \right] + \left(\frac{32768}{(3 \cdot 5 \cdot 17 \cdot 257)+1} - 1 \right)$$

$$M_p = \frac{2^{61}-1}{2305843009213693951}$$

$$\begin{aligned}
&= \left[\binom{2^{46}-1}{(2147483647)(32768)+(32768-1)} 70368744177663 \right] (32768) + (32768 - 1) \\
M_p &= \binom{2^{89}-1}{(2305843009213693951)(32768)\left(\frac{32768}{4}\right)} 618970019642690137449562111 \\
&= \left[\binom{2^{74}-1}{(2305843009213693951)\left(\frac{32768}{4}\right)} (18889465931478580854783) \right] (32768) + (32768 - 1) \\
M_p &= \binom{2^{107}-1}{(618970019642690137449562111)(32768)(8)} 162259276829213363391578010288127 \\
&= \left[\binom{2^{92}-1}{(618970019642690137449562111)(8)} (4951760157141521099596496895) \right] (32768) + (32768 - 1) \\
M_p &= \binom{2^{127}-1}{(162259276829213363391578010288127)(32768)(32)} 170141183460469231731687303715884105727 \\
&= \left[\binom{2^{112}-1}{(162259276829213363391578010288127)(32)} (5192296858534827628530496329220095) \right] (32768) + (32768 - 1)
\end{aligned}$$

In short, we can represent a Mersenne Prime $Mp_k > 32768$
 $Mp_k = 2^p - 1 = [(2^{p-15} - 1) (32768)] + (32768 - 1)$

This can be done with the rest of inradii (2,8,128) of the Pythagorean Triangles related to Fermat Primes, so we have:

$$\begin{aligned}
Mp_k &> 2 \\
Mp_k &= 2^p - 1 = [(2^{p-1} - 1) (2)] + (2 - 1)
\end{aligned}$$

$$\begin{aligned}
Mp_k &> 8 \\
Mp_k &= 2^p - 1 = [(2^{p-3} - 1) (8)] + (8 - 1)
\end{aligned}$$

$$\begin{aligned}
Mp_k &> 128 \\
Mp_k &= 2^p - 1 = [(2^{p-7} - 1) (128)] + (128 - 1)
\end{aligned}$$

$$\begin{aligned}
Mp_k &> 32768 \\
Mp_k &= 2^p - 1 = [(2^{p-15} - 1) (32768)] + (32768 - 1)
\end{aligned}$$

Note that (1,3,7,15) are the Mersenne Numbers (page 5, table 2).

Following, we can find some congruences:

$$\begin{aligned}
31 &\equiv (15 \cdot 2) \pmod{1} \\
127 &\equiv (15 \cdot 8) \pmod{7} \\
524287 &\equiv (15 \cdot 32768) \pmod{32767} \\
8191 &\equiv (1023 \cdot 8) \pmod{7} \\
131071 &\equiv (1023 \cdot 128) \pmod{127} \\
31 &\equiv (3 \cdot 8) \pmod{7} \\
131071 &\equiv (3 \cdot 32768) \pmod{32767}
\end{aligned}$$

A remark on the non infinity of Mersenne Primes and Perfect Numbers.

We know that $a + b = |\vec{a} \cdot \vec{b}| = \cos(\theta) \|\vec{a}\| \|\vec{b}\|$.

We can see that $\cos \theta \|a\| \|b\| - \frac{1}{\sqrt{2}} \|a\| \|b\| = 2n = F_k - 1$ for every $\begin{vmatrix} n & 1 \\ n+1 & F_k \end{vmatrix} = Mn_k$

Given the matrix and it's respective pythagorean triple

$$\underbrace{\begin{bmatrix} n & 1 \\ n+1 & F_k \end{bmatrix}}_{(a,b,c)} \rightarrow b - a = Mp_k$$

As every Perfect Number have the form $Sp_k \cdot Mp_k = Pf_k$, $\begin{cases} a = 2n + 1 \\ c = (2n + 1)n + (n + 1) = 2n^2 + 2n + 1 \\ b = c - 1 = 2n^2 + 2n \end{cases}$
 $Sp_k \cdot Mp_k = n^2 \cdot (b - a) = n^2 \cdot (2n^2 + 2n - 2n - 1) = n^2 \cdot (2n^2 - 1) = 2n^4 - n^2 = Pf_k$

(Sp_k = Superperfect Number, Mp_k = Mersenne Prime, Pf_k = Perfect Number)

$$\text{Superperfect Number} = \frac{\text{Perfect number}}{\text{Mersenne prime}} = \frac{2n^4 - n^2}{2n^2 - 1} = n^2 \quad \forall n \neq \left\{ -\frac{1}{\sqrt{2}} \text{ and } \frac{1}{\sqrt{2}} \right\}$$

Since the roots of $2n^2 - 1$ are $\left\{ -\frac{1}{\sqrt{2}} \text{ and } \frac{1}{\sqrt{2}} \right\}$, when $n^2 = Sp_k$ is confined to these limits.

Now we can represent the whole system of Mersenne Primes and Perfect Numbers this way:

Mersenne Prime = $2n^2 - 1$ = *Parallelogram*.

Perfect Number = $2n^4 - n^2$ = *Parallelepiped volume*.

SuperPerfect Number = n^2 = *Height*.

So, given 'n' we have:

$$n \Rightarrow \underbrace{\begin{bmatrix} n & 1 \\ n+1 & 2n+1 \end{bmatrix}}_{\text{Parallelogram}} \Rightarrow \underbrace{\begin{bmatrix} 1 & 2n+1 & n \\ n & n+1 & n^2 \\ n+1 & 1 & n \end{bmatrix}}_{\text{Parallelepiped volume}}$$

Example:

$$A = \begin{bmatrix} 2 & 1 \\ 3 & 5 \end{bmatrix} \Rightarrow |A| = 7 \Rightarrow A' = \begin{bmatrix} 1 & 5 & 2 \\ 2 & 3 & 4 \\ 3 & 1 & 2 \end{bmatrix} \Rightarrow |A'| = 28$$

$$A = \begin{bmatrix} 4 & 1 \\ 5 & 9 \end{bmatrix} \Rightarrow |A| = 31 \Rightarrow A' = \begin{bmatrix} 1 & 9 & 4 \\ 4 & 5 & 16 \\ 5 & 1 & 4 \end{bmatrix} \Rightarrow |A'| = 496$$

$$A = \begin{bmatrix} 8 & 1 \\ 9 & 17 \end{bmatrix} \Rightarrow |A| = 127 \Rightarrow A' = \begin{bmatrix} 1 & 17 & 8 \\ 8 & 9 & 64 \\ 9 & 1 & 8 \end{bmatrix} \Rightarrow |A'| = 8128$$

and so on.

1. Starting Point: The Parabola

Consider the quadratic function:

$$f(n) = 2n^2 - F_k$$

where F_k is the k -th Fermat number or Fermat-like number of the form:

$$F_k = 2^{2^k} + 1$$

The function $f(n)$ is symmetric about its vertex at:

$$n = \sqrt{\frac{F_k}{2}}$$

Let n_1 and n_2 be symmetric around this vertex, so that:

$$f(n_1) = -x, \quad f(n_2) = x$$

Then the values $|f(n_1)|$ and $|f(n_2)|$ are symmetric opposites.

2. Constructing the Pythagorean Triple

We define:

$$a = |f(n_1)|, \quad b = |f(n_2)|, \quad c = F_k$$

Then:

$$a^2 + b^2 = c^2$$

i.e., (a, b, c) is a **Pythagorean triple**.

3. The Matrix of Derivation

Given a primitive Pythagorean triple $[a, b, c]$, define:

$$\frac{c-a}{b} = \frac{q}{p}, \quad \frac{c-b}{a} = \frac{q'}{p'} \quad \Rightarrow \quad M = \begin{bmatrix} q & q' \\ p & p' \end{bmatrix}$$

It can be shown that:

$$p^2 + q^2 = c = p'^2 + q'^2$$

Thus, the triple is encoded in the matrix M , and its structure preserves the norm:

$$\|\vec{v}\|^2 = p^2 + q^2 = c$$

4. Recursive Fermat-like Construction

From the triple (a, b, c) , define:

$$F_{k+1} = 2^{2^k-1} \cdot a \cdot b + c$$

This recurrence generates the next Fermat-like number.


5. Summary

The recursive structure:

$$F_{k+1} = 2^{2^k-1} \cdot a_k \cdot b_k + F_k$$

emerges from selecting symmetric points on the parabola $f(n) = 2n^2 - F_k$, which produce the legs a_k and b_k of a right triangle with hypotenuse F_k

Figure 2: Points: $(1,-3), (2,3)$ and $(1,-15), (4,15)$



Parbola Distance between points.

$$\begin{aligned}
 f(n) &= 2n^2 - 5 \\
 d[1, -3]; [2, 3] \\
 &= \sqrt{37} \\
 &= \sqrt{1^2 + 6^2} \\
 &= \sqrt{1^2 + (2 \cdot 3)^2} \\
 &= 6.0827625302982 \approx 6 \\
 \text{Equation: } y &= 6x - 9, \frac{\Delta y}{\Delta x} = 6
 \end{aligned}$$

$$\begin{aligned}
 f(n) &= 2n^2 - 17 \\
 d[1, -15]; [4, 15] \\
 &= 3\sqrt{101} = 3\sqrt{1^2 + 10^2} \\
 &= 3\sqrt{1^2 + (2 \cdot 5)^2} \\
 &= 3\sqrt{1^2 + (2(1^2 + 2^2))^2} \\
 &= 30.149626863363 \approx 30 \\
 \text{Equation: } y &= 10x - 25, \frac{\Delta y}{\Delta x} = 10
 \end{aligned}$$

$$\begin{aligned}
 f(n) &= 2n^2 - 257 \\
 d[1, -255]; [16, 255] \\
 &= 3 \cdot 5\sqrt{1157} \\
 &= 3 \cdot 5\sqrt{1^2 + 34^2} \\
 &= 3 \cdot 5\sqrt{1^2 + (2 \cdot 17)^2} \\
 &= 3 \cdot 5\sqrt{1^2 + (2 \cdot (1^2 + 4^2))^2} \\
 &= 3 \cdot 5\sqrt{1^2 + (3^2 + 5^2)^2} \\
 &= 510.22054055085 \approx 510 \\
 \text{Equation: } y &= 34x - 289, \frac{\Delta y}{\Delta x} = 34
 \end{aligned}$$

$$\begin{aligned}
 f(n) &= 2n^2 - 65537 \\
 d[1, -65535]; [256, 65535] \\
 &= 3 \cdot 5 \cdot 17\sqrt{17 \cdot 1541} \\
 &= 3 \cdot 5 \cdot 17\sqrt{264197} \\
 &= 3 \cdot 5 \cdot 17\sqrt{264197} \\
 &= 3 \cdot 5 \cdot 17\sqrt{1^2 + 514^2} \\
 &= 3 \cdot 5 \cdot 17\sqrt{1^2 + (2 \cdot 257)^2} \\
 &= 3 \cdot 5 \cdot 17\sqrt{1^2 + (2(1^2 + 16^2))^2} \\
 &= 3 \cdot 5 \cdot 17\sqrt{1^2 + (15^2 + 17^2)^2} \\
 &= 131070.24805427 \approx 131070 \\
 \text{Equation: } y &= 514x - 66049, \frac{\Delta y}{\Delta x} = 514
 \end{aligned}$$

$$\begin{aligned}
& f(n) = 2n^2 - 4294967297 \\
& d[1, -4294967295]; [65536, 4294967295] \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 13 \sqrt{17 \cdot 5979949} \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 13 \sqrt{(1^2 + 4^2) \cdot (365^2 + 2418^2)} \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 13 \sqrt{(1^2 + 4^2) \cdot (14^2 + 15^2)^2 + 2418^2} \\
& = 8589934590.2501 \approx 8589934590 \\
& \text{Equation: } y = 131074x - 4295098369, \frac{\Delta y}{\Delta x} = 131074
\end{aligned}$$

$$\begin{aligned}
& f(n) = 2n^2 - 18446744073709551617 \\
& d[1, -18446744073709551615]; [4294967296, 18446744073709551615] \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \sqrt{316922321684979699947550670853} \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 65537 \sqrt{17 \cdot 941 \cdot 827009 \cdot 5577388969} \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 65537 \sqrt{17 \cdot (10^2 + 29^2)(545^2 + 728^2) \cdot (25688^2 + 70125^2)} \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \sqrt{562958543486978^2 + 65537^2} \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \sqrt{(2 \cdot 65537 \cdot 641 \cdot 6700417)^2 + 65537^2}
\end{aligned}$$

$$\begin{aligned}
& f(n) = 2n^2 - 340282366920938463463374607431768211457 \\
& d[1, -340282366920938463463374607431768211455]; \\
& [18446744073709551616, 340282366920938463463374607431768211455] \\
& = 714134895 \sqrt{908193463781113254077614364079150442524006982184239653426533} \\
& = 3 \cdot 5 \cdot 17^2 \cdot 257 \cdot 65537 \cdot 641 \cdot 6700417 \sqrt{13 \cdot 4166329117 \cdot 86957023859186942169125453} \\
& 2n^2 - 115792089237316195423570985008687907853269984665640564039457584007913129639937 \\
& d[1, -115792089237316195423570985008687907853269984665640564039457584007913129639935] \\
& [340282366920938463463374607431768211456, \\
& 115792089237316195423570985008687907853269984665640564039457584007913129639935] \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 65537 \cdot 641 \cdot 6700417 \cdot 274177 \cdot 67280421310721 \\
& \sqrt{17 \cdot 37 \cdot 6521 \cdot 4041028932361649884541 \cdot 27943581974981845864229861272202891692859495016013}
\end{aligned}$$

$$\begin{aligned}
& 2n^2 - 1340780792994259709957402499820584612747936582059239337772356144372176403007354697 \\
& 6801874298166903427690031858186486050853753882811946569946433649006084097 \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 65537 \cdot 641 \cdot 6700417 \cdot 274177 \cdot 67280421310721 \cdot 59649589127497217 \\
& \cdot 5704689200685129054721 \\
& \sqrt{17 \cdot 13 \cdot 692389 \cdot 5908249 \cdot 22559964396494555046199349 \cdot 10362691798570438439648634314849}
\end{aligned}$$

$$\begin{aligned}
& 2n^2 - 1797693134862315907729305190789024733617976978942306572734300811577326758055009631327084 \\
& 773224075360211201138798713933576587897688144166224928474306394741243777678934248654852763022 \\
& 196012460941194530829520850057688381506823424628814739131105408272371633505106845862982399472 \\
& 45938479716304835356329624224137217 \\
& = 3 \cdot 5 \cdot 17 \cdot 257 \cdot 65537 \cdot 641 \cdot 6700417 \cdot 274177 \cdot 67280421310721 \\
& \cdot 59649589127497217 \cdot 5704689200685129054721 \cdot 1238926361552897 \\
& \sqrt{17 \cdot 661 \cdot 22236089 \cdot 17301097129 \cdot 2434742448517 \cdot 264588417014506606729}
\end{aligned}$$

References

- [1] *H. Lee Price.*
The Pythagorean Tree: A New Species.
<https://arxiv.org/pdf/0809.4324.pdf>.
- [2] *University of San Antonio.*
The Geometric Interpretation of the Determinant.
https://mathresearch.utsa.edu/wiki/index.php?title=The_Geometric_Interpretation_of_the_Determinant
- [3] *On Primitive Pythagorean Triples*
<https://hal.science/hal-03247261/document>,page 8.
- [4] *George E. Andrews.*
Number Theory.
Chapter 7.Primitive Roots,p 97
Dover Publications, Inc. New York.1994.
- [5] *Notes on Perfect Numbers*
Geometric Representation of Perfect Numbers.p 15-16.
<https://hal.science/hal-01911641v7>
- [6] *University of Florida.Proof of the Cotes Theorem.*
<https://mae.ufl.edu/~uhk/COTES%20THEOREM.pdf>