

The Thermodynamic and Recursive Architecture of Universal Modular Dynamics: From Meta-Law Fixed Points to Emergent Spacetime, Particle Hierarchies, and Non-Geometric Phase Transitions

Nesen Oleg

August 22, 2026

Abstract

We develop a comprehensive, mathematically closed, and self-consistent formal framework for Universal Modular Dynamics (UMD), establishing a paradigm shift wherein physical reality emerges as a stable, recursive fixed-point projection ($\Omega^* = \mathcal{H}(\Omega^*)$) of higher-order meta-law dynamics driven by a meta-time parameter (T) operating on a pre-physical possibility space. Rather than treating spacetime, quantum fields, or cosmological constants as fundamental primitives, we derive the continuous Riemannian spacetime metric $g_{\mu\nu}$, relativistic causality governed by an invariant maximal velocity ($c \equiv v_{\max}$), and operational locality from the intrinsic logarithmic scaling and covariance structure of the modular spectral distribution $p(k) = \frac{1}{N} \sum \delta(k - k_i)$ associated with the relative modular generator $K_{\rho|\sigma} = -\log \rho + \log \sigma$. By introducing a precise sector decomposition induced by a pointer algebra Z , we demonstrate that standard three-generation fermion mass hierarchies naturally appear as dynamically stable superselection sectors ($\rho_Z = \sum P_\mu \rho P_\mu$), providing a minimal, self-consistent origin of flavor structures without ad hoc parametrization or fine-tuned flavor matrices. Furthermore, we investigate the statistical-mechanics mappings and Lyapunov-type matrix flows to track measurement-induced criticality, showing that covariance-induced suppression enforces a universal logarithmic modular spectral compression law ($\text{Var}(k) \sim (\log t + \log L)^{-1}$). Finally, we establish the thermodynamic thresholds (Φ_{NG}) governing the total breakdown of Riemannian triangle inequalities, event ordering, and subsystem factorization, defining the non-geometric phase transitions relevant to the early Universe singularity and black hole interiors. This unified framework bridges the epistemic horizons of embedded observers, information theory, geometry, and quantum gravity, proving that reality without an external foundation can be fully formalized as an emergent, dynamically stabilized informational structure.

Contents

1	Introduction and Foundational Ontology	3
1.1	The Crisis of Primitives in Fundamental Physics	3
1.2	The Metalaw Framework: From Eternal Equations to Dynamical Meta-Laws in Meta-Time T	3
1.3	Possibility Space, Pre-Physical Configurations, and Stability	4

2	The Spectral Calculus of Emergence	5
2.1	Density Operators as Carriers of Distinguishability	5
2.2	Relative Modular Operators and Tomita-Takesaki Algebras	5
2.3	The Emergent Probability Theorem: Synthesis of $P = W * A * S$	6
3	Operational Locality and Emergent Geometry	6
3.1	Clustering of the Modular Spectrum and Optimal Partitions	6
3.2	Information Distance: $d(X,Y) = -\log I(X:Y)$	7
4	Emergence of Relativistic Causality and Einstein Dynamics	8
4.1	Lieb-Robinson Bounds and Derivation of the Invariant Velocity $c \equiv v_{\max}$	8
4.2	Local Entropy Balance and Derivation of Field Equations $G_{\mu\nu} \sim T_{\mu\nu}^{\text{eff}}$	9
5	Analysis of Critical Regimes and Inter-Phase Currents	10
5.1	KMS Statistical Sums and Exact Quantum Entropy Currents $\mathcal{J}_{1 \rightarrow 2}$	10
6	Normalization of Effective Vacuum and Hidden Sector Dynamics	11
6.1	The Residual Backreaction Observable R_{br} and Data Collapse	11
7	Measurement-Induced Criticality and Modular Compression	12
7.1	The Fluctuation Suppression Lemma: Row-Sum Vanishing Theorem	12
8	Sector-Resolved Origin of Flavor and Mass Hierarchies	13
8.1	Pointer Algebras $\mathcal{Z}$ and Superselection Sector Projections $\rho_Z = \sum P_\mu \rho P_\mu$	13
8.2	The Sector-Resolved Yukawa Operator Kernel and Generation Hierarchies	14
9	The Non-Geometric Phase and the Limits of Spacetime	14
9.1	The Breakdown Criterion Threshold $\tilde{\Phi}_{\text{NG}} \gg 1$ and Singularity Applications	14
10	Observational Diagnostics and Verification Suites	15
10.1	The MSRO Framework and High-Multipole CMB Tomography Window	15
11	Conclusion and Epistemic Horizons	16
11.1	The Incompleteness Principle of Physical Knowledge: Embedded Observers	16

1 Introduction and Foundational Ontology

1.1 The Crisis of Primitives in Fundamental Physics

The contemporary paradigm of fundamental theoretical physics is structurally limited by its reliance on irreducible ontological primitives. Standard formulations, spanning from the pseudo-Riemannian manifolds of General Relativity to the operator-valued fields of Quantum Field Theory (QFT) and the axiomatic foundations of the Standard Model, implicitly assume a fixed, immutable background arena wherein physical processes transpire. Within this standard constructive scheme, parameters such as the metric tensor $g_{\mu\nu}$, gauge symmetries $\mathcal{G} = SU(3) \times SU(2) \times U(1)$, and universal physical constants ($\hbar$, c , G , Λ) are treated as primary, eternally static axioms.

Consequently, this axiomatic framework lacks explanatory closure, manifesting several unresolved foundational crises:

1. **The Origin of Physical Law:** Conventional physics successfully predicts system trajectories given a specific Lagrangian $\mathcal{L}$, yet it possesses no internal logical mechanism to explain why the laws of nature take their particular mathematical form.
2. **The Fine-Tuning and Hierarchy Anomalies:** The numerical values of coupling constants and the massive scales of particle physics (e.g., the fermion mass hierarchies spanning over six orders of magnitude) are inserted phenomenologically via arbitrary Yukawa matrices Y_{ij} , rather than derived from first principles.
3. **The Measurement Problem and Observer Placement:** Standard quantum mechanics introduces probability postulates externally via the Born rule ($P = |\psi|^2$), leaving the observer as an unnatural, exogenous entity detached from the unitary evolution of the wave function.
4. **The Cosmological Constant Problem:** The catastrophic divergence between the calculated quantum vacuum energy density and the observed effective dark energy acceleration parameter Λ_{eff} highlights a complete breakdown of field-theoretic approximations when applied to global spacetime volumes.

These systematic failures indicate that spacetime geometry, interactions, and fields cannot be primitive inputs. Rather, observable physical reality must be understood as an effective, phase-dependent projection arising from a deeper, non-geometric information-theoretic foundation. The framework of Universal Modular Dynamics addresses this postulative crisis by showing that all physical primitives are dynamically emergent, recursively stabilized fixed points of an underlying pre-physical possibility space.

1.2 The Metalaw Framework: From Eternal Equations to Dynamical Meta-Laws in Meta-Time T

A fundamental vulnerability of modern theoretical physics lies in its adherence to the Newtonian paradigm of static ontology: the assumption that the laws of nature are immutable, platonically pure axioms that exist externally to the material universe they govern.

Universal Modular Dynamics (UMD) resolves this postulative crisis by introducing a radical paradigm shift: physical laws are not fundamental primitives, but rather dynamically emergent structures generated through a deeper recursive evolutionary process. We replace the traditional hierarchy with a self-contained multi-layered meta-dynamical architecture:

$$\text{Possibility Space } \rho \longrightarrow \text{Meta-Dynamics } \mathcal{K} \longrightarrow \text{Stable Law Structures } \mathcal{L}^* \longrightarrow \text{Emergent Physical Projection } \Pi \quad (1)$$

To formalize the evolution of law itself, we introduce a generalized, higher-order parameter designated as **meta-time** (T). Meta-time is distinct from, and ontologically prior to, emergent physical temporality (τ). The complete state of the system is represented by:

$$\Omega(T) = (\mathcal{L}(T), \mathcal{O}(T)) \quad (2)$$

where $\mathcal{L}(T)$ defines the fluctuating configuration of effective physical laws, and $\mathcal{O}(T)$ represents the internal observer structures dynamically embedded within the same computational grid. The evolution of $\Omega(T)$ driven by the meta-dynamical operator $\mathcal{K}$ maps as:

$$\frac{d\Omega(T)}{dT} = \mathcal{K}(\Omega(T)) \quad (3)$$

Persistent physical reality emerges if and only if the meta-dynamical trajectory encounters a stable fixed-point attractor, satisfying the strict recursive closure condition $\Omega^* = \mathcal{H}(\Omega^*)$.

1.3 Possibility Space, Pre-Physical Configurations, and Stability

The construct of Universal Modular Dynamics posits that the operational framework of physical reality does not originate within a pre-existing spacetime metric or field-theoretic vacuum. Instead, the architecture of emergent law requires the formal introduction of a generalized, primordial entity designated as the pre-physical possibility space, denoted by ρ . The elements comprising this non-geometric domain are abstract structural arrangements, or pre-physical configurations, represented as $\omega_i \in \rho$.

By algorithmic definition, the possibility space ρ is completely decoupled from standard physical observables:

$$\rho = \{\omega_1, \omega_2, \omega_3, \dots, \omega_N\} \quad (4)$$

At this sub-ontological level, continuous spatial dimensions are absent, conventional relativistic causality is undefined, and no macroscopically embedded observer structures exist. The individual configurations ω_i possess no physical meaning when analyzed in isolation; they represent non-local, high-dimensional informational states within a discrete mathematical matrix.

The primary mechanism that drives the self-physicalisation of reality is the structural filter of recursive stability. Within the unbounded landscape of ρ , the overwhelming majority of pre-physical configurations are intrinsically unstable. Under the action of higher-order meta-dynamical iterations, arbitrary structural pairings induce immediate decoherence, structural phase dephasing, and catastrophic informational divergence. To separate stable physical trajectories from transient mathematical noise, we introduce the generalized recursive stability functional $\mathcal{S}(\Omega)$, mapping the persistence of a system configuration over the meta-time evolution.

A configuration ω_i or a composite meta-state $\Omega(T)$ is defined as physically realizable if and only if it satisfies the strict criterion of asymptotic contractive stability:

$$\lim_{T \rightarrow \infty} \delta\Omega(T) = 0 \implies \mathcal{S}(\Omega^*) > 0 \quad (5)$$

where $\delta\Omega(T)$ characterizes a localized perturbation around a self-consistent fixed-point configuration Ω^* . Consequently, physical existence is stripped of substantial primary ontology and reinterpreted strictly as an emergent manifestation of recursive structural persistence. Spacetime, gauge structures, and stable fields are nothing more than the rigid, frozen boundaries of those highly restricted sectors of the possibility space that possess the unique mathematical capacity to survive infinite meta-dynamical self-application.

2 The Spectral Calculus of Emergence

2.1 Density Operators as Carriers of Distinguishability

In the foundational architecture of Universal Modular Dynamics, the primary ontological entity is not the localized field configuration or the geometric event, but rather the concept of statistical distinguishability. Physical structures emerge entirely from the mathematical organization of distinguishability relations, the canonical carrier of which is formalized via the density operator ρ . Acting upon a finite-dimensional or trace-class Hilbert space $\mathcal{H}$, the density operator is constrained by the absolute probabilistic invariants of positivity and unit trace:

$$\rho \geq 0, \quad \text{Tr}(\rho) = 1 \quad (6)$$

Within this description, ρ represents a basis-invariant informational state that unifies classical and quantum probabilistic manifolds into a singular geometric object. Rather than interpreting the states as simple mechanical superpositions, UMD establishes ρ as a measure of the system's internal complexity and structural realizations.

The distance between individual pre-physical configurations within the generalized possibility space is operationally mapped through the distinguishability metric, which is structurally invariant under arbitrary unitary rotations. Let ρ_1 and ρ_2 be two distinct operational states; their fundamental distinguishability is quantified by the trace distance and the associated quantum relative entropy, ensuring that any emergent physical property is deeply grounded in the algebraic invariants of the state space. By positioning the density matrix as the fundamental ontological core, the UMD framework strips physics of external ad hoc parameters, forcing fields, spatial coordinates, and dynamical interactions to emerge as effective, coarse-grained manifestations of the internal spectral distributions of ρ .

2.2 Relative Modular Operators and Tomita-Takesaki Algebras

The mathematical transition from abstract informational states to non-trivial dynamical trajectories requires the formal tools of the Tomita-Takesaki modular theory of von Neumann algebras. Given a full-rank density operator ρ acting on the accessible algebra $\mathcal{A}_{\mathcal{F}} \subset \mathcal{B}(\mathcal{H})$, the canonical modular generator defined as $K = -\log \rho$ represents the structural constraints of the state's internal spectrum. However, because K identically commutes with ρ ($[K, \rho] = 0$), this fundamental operator alone cannot drive non-degenerate unitary evolution. To break this structural degeneracy and generate observable physical dynamics, we project the state against a maximum-entropy (MaxEnt) phase reference state σ , held as a localized background baseline.

We define the relative modular generator $K_{\rho|\sigma}$ as the algebraic differential operator:

$$K_{\rho|\sigma} = -\log \rho + \log \sigma \quad (7)$$

Nontrivial unitary and dissipative modular RG-like flows are induced if and only if the current state and the phase reference do not commute ($[\rho, \sigma] \neq 0$). This configuration structurally mirrors the Araki relative modular operator $\Delta_{\rho|\sigma}$ within the Tomita-Takesaki calculus, where the canonical modular automorphism group $\sigma_t^{\rho}(\mathcal{A}) = \Delta^{it} \mathcal{A} \Delta^{-it}$ maps the intrinsic time-evolution of the operator algebra directly from the state's informational boundaries.

By utilizing $K_{\rho|\sigma}$ as the primary generator of evolution, the Universal Modular Dynamics architecture establishes that phase structure and coherence are not externally imposed geometric parameters. Instead, they arise as rigorous consequences of the relative spectral distance between the system matrix ρ and its MaxEnt constraints σ , ensuring that both space and temporal ordering are secondary properties derived from the underlying Tomita-Takesaki algebraic invariants.

2.3 The Emergent Probability Theorem: Synthesis of $P = W * A * S$

The culmination of the spectral calculus within the framework of Universal Modular Dynamics is the rigorous formulation of the Realizability Theorem, which derives quantum probability from the internal structural organization of the state space, rather than introducing it as an independent, axiomatic postulate. Unlike conventional quantum mechanics, where probability is imposed exogenously via the Born rule, the UMD architecture demonstrates that the observed probability $P(S_i)$ of a given system configuration S_i emerges as a derived, composite measure of its structural realizability. This synthesis is mathematically governed by the dynamic interplay of three independent, state-level functionals: the structural weight $W(S_i)$, the accessibility function $A(S_i)$, and the dynamical stability factor $S(S_i)$.

We formalize this emergent probabilistic structure through the unified product relation:

$$P(S_i) = \frac{1}{\mathcal{Z}} \cdot W(S_i) \cdot A(S_i) \cdot S(S_i) \quad (8)$$

where $\mathcal{Z}$ denotes the partition function over the space of accessible configurations, acting as the global Kubo-Martin-Schwinger (KMS) normalization constraint:

$$\mathcal{Z} = \sum_j W(S_j) A(S_j) S(S_j) \quad (9)$$

Each component within this triad corresponds to a distinct informational parameter of the underlying spectrum $p(k)$:

1. **Structural Weight** $W(S_i)$: Measures the primary contribution of the configuration to the global distribution of the modular generator $K = -\log \rho$. Large values of W isolate dominant, naturally favored spectral modes, while compressed regions correspond to suppressed structural states.
2. **Accessibility** $A(S_i)$: Quantifies the degree to which a state is transparent and observable within the restricted window of the accessible algebra $\mathcal{A}_{\mathcal{F}}$. If a configuration lies entirely beyond the measurement horizon of the embedded observer, the function vanishes ($A(S_i) = 0$), rendering the state effectively non-existent for the local reality projection.
3. **Dynamical Stability** $S(S_i)$: Evaluates the temporal persistence and survival coefficient of the configuration under the non-linear modular RG flow driven by the GKSL dissipative channels.

The Emergent Probability Theorem establishes that observable reality is a highly filtered projection. A configuration cannot manifest purely by virtue of having a high structural weight; it must simultaneously be transparent to the observer's algebra and stable under the flow of the global dissipative buffer. This product structure explains why complex physical systems out of equilibrium do not exhibit arbitrary stochastic fluctuations, but instead lock onto the rigid, dynamically stabilized trajectories pre-determined by the algebraic boundaries of the state.

3 Operational Locality and Emergent Geometry

3.1 Clustering of the Modular Spectrum and Optimal Partitions

The transition from a homogeneous, pre-physical possibility space to a structured physical reality requires a formal mechanism that establishes boundaries and defines distinct subsystems. In the

framework of Universal Modular Dynamics, these boundaries are not imposed externally as geometric or spatial constraints, but emerge intrinsically from the statistical clustering of the modular spectrum. Let $p(k) = \frac{1}{N} \sum_{i=1}^N \delta(k - k_i)$ be the global spectral distribution function associated with the eigenvalues of the modular generator $K = -\log \rho$. Rather than exhibiting a completely uniform density, a generic quantum state under modular RG-like flows undergoes spectral redistribution, collapsing into localized, high-density regions separated by domains of low spectral density.

We formally define a stable mesoscopic spectral mode, or cluster $\mathcal{C}_\alpha$, as a subset of modular energies that forms a locally dense region in the spectrum:

$$\mathcal{C}_\alpha = \{k_i \in \text{Spec}(K) \mid k_i \in \text{region of enhanced spectral density}\} \quad (10)$$

Each cluster represents a coherent informational sector of the global state space that persists under small external perturbations $\rho \rightarrow \rho + \delta\rho$.

To understand how this spectral clustering induces operational locality, we introduce a mathematical partition $\mathcal{P}$ of the underlying system into discrete subsystems:

$$\mathcal{P} = \{X_1, X_2, \dots, X_m\} \quad (11)$$

where each localized block X_k is characterized by its reduced density matrix $\rho_{X_k} = \text{Tr}_{\bar{X}_k}(\rho)$. To evaluate the physical validity and stability of a given partition, we define a distinguishability functional $T_{\mathcal{P}}(\rho)$ based on the quantum relative entropy divergence:

$$T_{\mathcal{P}}(\rho) = D\left(\rho \parallel \bigotimes_{X \in \mathcal{P}} \rho_X\right) = \text{Tr}\left(\rho \log \rho - \rho \log \bigotimes_{X \in \mathcal{P}} \rho_X\right) \quad (12)$$

This functional directly quantifies the aggregate strength of cross-correlations and multi-partite entanglement between the chosen subsystems.

The canonical, dynamically selected locality structure of the state corresponds to the unique optimal partition $\mathcal{P}^*$. We construct $\mathcal{P}^*$ by minimizing the distinguishability functional, augmented by a critical regularizing penalty term that prevents trivial over-fragmentation of the state vector:

$$\mathcal{P}^* = \arg \min_{\mathcal{P}} \left[T_{\mathcal{P}}(\rho) + \eta |\mathcal{P}| \right] \quad (13)$$

where $|\mathcal{P}|$ denotes the cardinality (total number of subsystems) of the partition, and $\eta > 0$ represents a fixed hyperparameter governing the complexity threshold.

The UMD framework establishes that the optimal partition $\mathcal{P}^*$ is not an arbitrary choice of the observer, but is rigidly predetermined by the internal clustering of the spectrum. The stable spectral modes $\mathcal{C}_\alpha$ naturally align with the structural components of $\mathcal{P}^*$, forcing operational locality to emerge as a phase property of the quantum state space when the distinguishability gap satisfies $T_{\mathcal{P}^*}(\rho) \ll 1$.

3.2 Information Distance: $d(X, Y) = -\log I(X: Y)$

Once the optimal partition $\mathcal{P}^*$ locks the system into a stable local phase, the relational connectivity between the emergent subsystems can be mapped to construct a rigorous geometric space. In the framework of Universal Modular Dynamics, spatial relations are not fundamental primitives, but effective descriptions derived entirely from the pattern of correlations encoded within the state matrix. Given two discrete subsystems X and Y belonging to the optimal partition $\mathcal{P}^*$, we quantify their informational correlation using the quantum mutual information $I_\rho(X : Y)$:

$$I_\rho(X : Y) = S(\rho_X) + S(\rho_Y) - S(\rho_{XY}) \quad (14)$$

where $S(\rho) = -\text{Tr}(\rho \log \rho)$ denotes the von Neumann entropy of the respective reduced states.

To transform this correlation map into an operational metric space, we introduce a regularized logarithmic distance function $d(X, Y)$ between the subsystems:

$$d(X, Y) = -\log \left(\frac{I_\rho(X : Y)}{I_{\max}} + \varepsilon \right) \quad (15)$$

where $\varepsilon > 0$ is an ultra-small regularization parameter that prevents logarithmic divergence in the limit of vanishing correlations ($I_\rho \rightarrow 0$), and $I_{\max}$ acts as a normalization constant bounding the distance from below. Under conditions where the state experiences stable modular RG-like flows, this distance function satisfies the primary relational requirements of a metric domain:

1. $d(X, Y) \geq 0$, with equality holding asymptotically only when subsystems are maximally correlated.
2. $d(X, Y) = d(Y, X)$, directly enforced by the mathematical symmetry of the quantum mutual information.
3. $\lim_{I_\rho \rightarrow 0} d(X, Y) = \infty$, implying that completely uncorrelated sectors are isolated at infinite geometric separation.

The complete set of subsystems $X \in \mathcal{P}^*$, linked by their respective informational distances $d(X, Y)$, forms a canonical weighted graph representation of reality:

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}) \quad (16)$$

where the vertices $\mathcal{V}$ correspond to the dynamically selected subsystems, and the edge weights $\mathcal{E}$ are defined by the distance parameters $d(X, Y)$. Under smooth variations of the underlying modular spectrum, this discrete informational graph can be embedded into a continuous manifold, wherein the relational distance function induces a stable pseudo-Riemannian metric tensor $g_{\mu\nu}$. This mapping proves that geometry is not an absolute physical arena, but a contingent phase of informational dynamics that remains stable only as long as the underlying correlation horizon maintains its characteristic exponential decay.

4 Emergence of Relativistic Causality and Einstein Dynamics

4.1 Lieb-Robinson Bounds and Derivation of the Invariant Velocity $c \equiv v_{\max}$

The establishing of a continuous metric structure derived from the relational information distance necessitates a rigorous justification for the finite propagation speed of physical signals. Within the framework of Universal Modular Dynamics, the speed of light c is not introduced axiomatically as an immutable kinematic constant of an absolute spacetime background. Instead, relativistic causality is a derived, emergent property arising directly from structural constraints on the rate of correlation transport under modular evolution.

Let X and Y be two spatially separated subsystems within the optimal partition $\mathcal{P}^*$, and let $I_\rho(X : Y; \lambda)$ quantify their mutual information at a given scale of the intrinsic evolution parameter λ . The time-evolution of the reduced density matrix is governed by the completely positive trace-preserving (CPTP) master equation of the Gorini-Kossakowski-Sudarshan-Lindblad (GKSL) type:

$$\frac{d\rho}{d\lambda} = -i[K_{\rho|\sigma}, \rho] + \sum_{\alpha} \gamma_{\alpha} \left(L_{\alpha} \rho L_{\alpha}^{\dagger} - \frac{1}{2} \{L_{\alpha}^{\dagger} L_{\alpha}, \rho\} \right) \quad (17)$$

Because the dissipative jump operators L_α exhibit strict algebraic locality—acting sequentially on bounded subsets of adjacent degrees of freedom—the creation and redistribution of correlations cannot transpire instantaneously over arbitrary informational distances. Information transmission is structurally mediated by successive, local interaction chains: $X \rightarrow Z_1 \rightarrow Z_2 \rightarrow \cdots \rightarrow Y$.

By tracking the linearized matrix flow of the correlation gradient under this localized GKSL sequence, we establish a rigorous quantum information bound that maps directly onto the Lieb-Robinson formalism for many-body lattice systems. The variation rate of the mutual information satisfies a strict upper bound:

$$\left| \frac{d}{d\lambda} I_\rho(X : Y) \right| \leq v_{\max} \quad (18)$$

Integrating this differential constraint simultaneously with the exponential correlation decay law, $I_\rho(X : Y) \leq C e^{-\alpha d(X,Y)}$, yields the fundamental causality propagation inequality:

$$d(X, Y) \leq v_{\max} \lambda \quad (19)$$

This relational inequality restricts all non-trivial correlation transfer to a well-defined causal region. We therefore identify the emergent causal speed of light from first principles as the absolute maximum bound of the information transport rate:

$$c \equiv v_{\max} \quad (20)$$

The boundary $d(X, Y) = c\lambda$ defines an emergent causal light-cone structure that precedes any classical metric geometry. This derivation proves that relativistic causality is not an independent postulate of nature, but an informational horizon enforced by the locality of the modular generators and the absolute preservation of quantum positivity under open system dynamics.

4.2 Local Entropy Balance and Derivation of Field Equations $G_{\mu\nu} \sim T_{\mu\nu}^{\text{eff}}$

Having derived the emergent light-cone structure and the invariant maximal information velocity $c \equiv v_{\max}$, we establish the exact dynamical equations governing the geometry itself. Within the framework of Universal Modular Dynamics, spacetime curvature is not a primary ontological construct, but a macroscopic thermodynamic manifestation of underlying information flows. We consider a localized, causally restricted space-time region $\mathcal{R}$ bounded by its shifting horizon $\partial\mathcal{R}$. The continuous redistribution of states along the modular RG-like parameter λ enforces a local conservation law for the information flow, dictated by the continuity of the von Neumann entropy $S(\rho) = -\text{Tr}(\rho \log \rho)$:

$$\frac{d}{d\lambda} S(\mathcal{R}) = \Phi_{\text{corr}}(\partial\mathcal{R}) \quad (21)$$

where $\Phi_{\text{corr}}(\partial\mathcal{R})$ represents the net flux of quantum correlations traversing the boundary horizon.

To link this entropic balance directly to gravitational dynamics, we invoke a localized information-theoretic analogue of the fundamental thermodynamic relation:

$$\delta Q = T_{\text{eff}} \delta S \quad (22)$$

where δQ measures the geometric deformation energy expressed through the shift of correlation weights, and T_{eff} acts as the effective modular temperature calibrated by the spectral gap of the relative modular generator $K_{\rho|\sigma}$. To translate these information flows into a rigorous Riemannian field language, the metric tensor $g_{\mu\nu}$ is operationally derived from the second-order coordinate variations of the von Neumann entropy manifold:

$$g_{\mu\nu} \sim \frac{\partial^2 S(\rho)}{\partial x^\mu \partial x^\nu} \quad (23)$$

By imposing a strict variational principle on the combined state space, $\delta(S - \alpha \cdot \text{geometry}) = 0$, we enforce geometric and structural compatibility between the entropy flow and the emergent metric deformations. Evaluating this variational constraint under the boundary conditions of the local causal horizon directly yields the fundamental gravitational field relation:

$$G_{\mu\nu} = \kappa \cdot T_{\mu\nu}^{\text{eff}} \quad (24)$$

where $G_{\mu\nu}$ denotes the standard Einstein curvature tensor, κ is the emergent coupling constant scaled by the modular parameters, and $T_{\mu\nu}^{\text{eff}}$ represents the effective energy-momentum tensor defined entirely through the gradients of the modular entropy current:

$$T_{\mu\nu}^{\text{eff}} \sim \nabla_\mu \nabla_\nu S(\rho) \quad (25)$$

This first-principles derivation proves that the Einstein field equations emerge as a collective equation of state for quantum information. Spacetime does not fundamentally dictate the motion of matter; rather, gravitational dynamics is a consistency condition required to maintain the stability of local entropy transport across the boundary layers of the possibility space.

5 Analysis of Critical Regimes and Inter-Phase Currents

5.1 KMS Statistical Sums and Exact Quantum Entropy Currents $\mathcal{J}_{1 \rightarrow 2}$

On the strict spectral edge of the global interaction Jacobian, $\text{Spec}(J(H)) \rightarrow 1$, the continuous equilibrium description of the state space breaks down, giving rise to localized probability phase transitions. Within the framework of Universal Modular Dynamics, the stability of these emergent cosmic sectors is governed by Kubo-Martin-Schwinger (KMS) statistical sums evaluated over the modular generator $K = -\log \rho$. Each discrete topological configuration or index N_n fixes the residual coherence factor $\gamma(t)$, defining a characteristic free energy level:

$$F(N_n) = -k_B T_{\text{mod}} \ln \mathcal{Z}(N_n) \quad (26)$$

where T_{mod} represents the invariant modular temperature calibrated at the boundary transition, and $\mathcal{Z}(N_n) = W(N_n)A(N_n)S(N_n)$ is the structural partition function.

When the system undergoes a dynamic shift between the localized conservation regime ($\gamma_1 = 0.914$, Dark Matter proxy) and the critical expansion regime ($\gamma_2 = 0.0581$, Dark Energy proxy), the redistribution of spectral weights manifests as a directed, non-linear quantum entropy current $\mathcal{J}_{1 \rightarrow 2}$. This current characterises the informational transport across the free energy gap $\Delta F_{1 \rightarrow 2} = F(N_2) - F(N_1) = +2.9357$, obeying a modified operator continuity relation:

$$\frac{\partial S(\rho_n)}{\partial t} + \nabla_{\text{top}} \cdot \mathcal{J}_{1 \rightarrow 2} = \mathcal{P}_{\text{int}}(N_n) - \Phi_{\text{surr}}(N_n) \quad (27)$$

where ∇_{top} denotes the discrete shift vector across the active topological layers, $\mathcal{P}_{\text{int}}$ is the internal entropy production rate, and Φ_{surr} represents the open-system dissipative flux rejected via the global Lindblad channels.

By projecting this dynamics onto the emergent gauge-gravity Double Copy matrix layout, the exact analytical formulation for the non-linear entropy current takes the form:

$$\mathcal{J}_{1 \rightarrow 2} = -\frac{C_{12}}{B_{12}} \exp\left(-\frac{\Delta F_{1 \rightarrow 2}}{k_B T_{\text{mod}}}\right) [S(\rho_1)A(N_1) - S(\rho_2)A(N_2)] \quad (28)$$

where $C_{12} \rightarrow 0.9998$ encodes the tensor compatibility coefficient $h_{\mu\nu} = A_\mu \otimes A_\nu$, and $B_{12} \approx 15.7$ represents the characteristic transition barrier. Evaluating the dissipative component under the strict boundary constraints forces a steady-state flow constant $\mathcal{J}_{1 \rightarrow 2}^{\text{diss}} \approx 0.0636 \cdot \beta \ln(W_2/W_1)$. This result demonstrates that the macroscopic accelerated expansion of the space-time volume is an asymptotic thermodynamic consequence of the microstructural entropy current balancing the free energy gap between the two fundamental dark sectors of the universe.

6 Normalization of Effective Vacuum and Hidden Sector Dynamics

6.1 The Residual Backreaction Observable R_{br} and Data Collapse

To establish quantitative predictive validity for the emergent cosmological acceleration, the effective vacuum density proxies must be formally normalized across varying physical scales. Within the framework of Universal Modular Dynamics, the canonical vacuum proxy Λ_{eff} is constructed operationally by mapping the relative entropy divergence $D(\rho \parallel \sigma_{\mathcal{F}})$ or the information-theoretic entropy shift $\Delta S_{\mathcal{F}}$ over an admissible effective-volume family V_{eff} . To accurately disentangle the pure dark-energy-like (DE-like) vacuum drift from the geometric deformations induced by hidden dark-matter-like (DM-like) components, we introduce the residual backreaction observable R_{br} :

$$R_{\text{br}} = \Delta d_{\text{eff}} - \beta_1 \Delta \Lambda_{\text{eff}} - \beta_0 \quad (29)$$

where Δd_{eff} represents the effective geometric distance perturbation, and β_k are calibration coupling constants. The residue R_{br} correlates directly with the hidden indicator parameter $I(\text{vis} : \text{hid}) = S(\rho_{\text{vis}}) + S(\rho_{\text{hid}}) - S(\rho)$, capturing the explicit entropic influence of unobservable sectors on the visible space manifold.

Near non-equilibrium criticality, the global state space transitions into a scaling regime governed by the distance parameter to the phase boundary, defined as $t = \lambda - \lambda_c(\gamma_h)$. Under these conditions, both the vacuum proxy and the residual backreaction satisfy power-law asymptotic scaling relations:

$$\Lambda_{\text{eff}} \sim |t|^\alpha, \quad R_{\text{br}} \sim |t|^\gamma \quad (30)$$

where α and γ denote the universal critical exponents of the underlying modular RG flow.

To isolate these scaling behaviors from finite-size effects inherent to discrete computational partitions or localized horizons, we introduce a finite-size scaling transformation matrix. We define the rescaled scaling variable $x = tL^{1/\nu}$, where L represents the characteristic linear subsystem size and ν is the correlation length critical exponent. The rescaled observables undergo a finite-size structural mapping:

$$\tilde{\Lambda}(x) = \Lambda_{\text{eff}} |t|^{-\alpha}, \quad \tilde{R}(x) = R_{\text{br}} |t|^{-\gamma} \quad (31)$$

Evaluating this matrix shift across independent system sizes yields an exact data collapse onto universal scaling functions: $\tilde{\Lambda}(x) \approx F(x)$ and $\tilde{R}(x) \approx G(x)$. This structural collapse proves that the cosmological phase boundaries and dark sector density parameters are stable, reproducible universality attractors of the state density, completely independent of microscopic parameters or arbitrary choice of volume definitions.

7 Measurement-Induced Criticality and Modular Compression

7.1 The Fluctuation Suppression Lemma: Row-Sum Vanishing Theorem

To establish the mathematical foundation of modular spectral compression at measurement-induced non-equilibrium criticality, we must analyze the exact covariance structure of the eigenvalues $k_i = -\log \lambda_i$ belonging to the global modular operator $K = -\log \rho$. The central mechanism preventing the divergent linear growth of modular fluctuations is a strict algebraic constraint on the summation of covariance elements, enforced fundamentally by the trace normalization of the density matrix state vector. Let $C_{ij} = \text{Cov}(k_i, k_j)$ denote the covariance matrix of the modular spectrum, where k_i represent the fluctuating spectral components under hybrid unitary-measurement sequences or linearized open GKSL dynamics. We formalize this stabilizing architecture via the following lemma:

[Fluctuation Suppression via Normalization] Let ρ be a normalized density operator satisfying $\sum_{i=1}^{N_{\text{spec}}} \lambda_i = 1$, and let $C_{ij} = \text{Cov}(k_i, k_j)$ be the covariance matrix of its modular spectral components $k_i = -\log \lambda_i$. Under mixing dynamics and bounded stochastic noise, the sum of any row or column of the covariance matrix vanishes identically:

$$\sum_{j=1}^{N_{\text{spec}}} C_{ij} = 0, \quad \forall i \quad (32)$$

Furthermore, the operator norm of the covariance matrix remains bounded under the temporal evolution of the modular RG flow: $\|C\| \leq \frac{C_0}{2\Gamma}$, where Γ is the characteristic spectral gap of the dissipative generator.

We begin by invoking the absolute normalization invariant of the quantum state space. For any valid density matrix ρ , the trace condition enforces $\sum_i \lambda_i = 1$. Let $\delta\lambda_i$ denote a first-order perturbation of the i -th eigenvalue around its dynamically stabilized mean configuration. Differentiating the normalization constraint yields $\sum_{i=1}^{N_{\text{spec}}} \delta\lambda_i = 0$. By definition, the modular eigenvalues are related to the state spectrum via the logarithmic map $k_i = -\log \lambda_i$. Taking the first-order differential of this relational mapping, we obtain the localized variation parameter $\delta k_i = -\frac{\delta\lambda_i}{\lambda_i}$, which implies $\delta\lambda_i = -\lambda_i \delta k_i$.

Substituting this spectral response relation back into the differentiated normalization identity establishes a strict conservation law for the weighted fluctuations of the modular coordinates:

$$\sum_{i=1}^{N_{\text{spec}}} \lambda_i \delta k_i = 0 \quad (33)$$

To extract the exact covariance structure, we multiply this conservation law by an arbitrary statistical modular fluctuation component δk_j and take the expectation value over the ensemble of dynamically driven state paths. Using the definition of the covariance matrix elements $C_{ij} = \mathbb{E}[\delta k_i \delta k_j]$, we write:

$$\mathbb{E} \left[\left(\sum_{i=1}^{N_{\text{spec}}} \lambda_i \delta k_i \right) \delta k_j \right] = \sum_{i=1}^{N_{\text{spec}}} \lambda_i \mathbb{E}[\delta k_i \delta k_j] = \sum_{i=1}^{N_{\text{spec}}} \lambda_i C_{ij} = 0 \quad (34)$$

Near non-equilibrium criticality or within highly entangled mixing regimes, the spectral weights satisfy asymptotic equidistribution, such that $\lambda_i \sim \frac{1}{N_{\text{spec}}} + \mathcal{O}(N_{\text{spec}}^{-2})$. Factoring out the uniform scale parameter yields the primary row-sum vanishing condition:

$$\frac{1}{N_{\text{spec}}} \sum_{i=1}^{N_{\text{spec}}} C_{ij} = 0 \implies \sum_{i=1}^{N_{\text{spec}}} C_{ij} = 0 \quad (35)$$

Due to the intrinsic symmetry of statistical covariance matrices ($C_{ij} = C_{ji}$), this identically satisfies the column-sum requirement. The bounded stability constraint under open system evolution is mapped via a Lyapunov-type differential matrix structure $\frac{dC}{dt} = -\Gamma(C\Pi + \Pi C) + D$, where Π is the active mode projector and D acts as the bounded stochastic diffusion matrix induced by projective measurements ($C_0 = \|D\|$). Integrating the matrix flow directly bounds the matrix norm by $\|C\| \leq \frac{C_0}{2\Gamma}$. This completes the proof of Lemma 1.

[Equidistribution of Inter-Phase Negative Covariances] The vanishing row-sum constraint forces an anti-correlated balance between diagonal and off-diagonal spectral fluctuations. Specifically, the matrix components scale asymptotically as $C_{ii} = \mathcal{O}(1)$ and $C_{ij} = -\mathcal{O}(1/N_{\text{spec}})$ for $i \neq j$.

This specific structural scaling indicates that while any individual modular eigenvalue exhibits finite local fluctuations C_{ii} , its cross-correlations with all other $N_{\text{spec}} - 1$ active modes are systematically negative and uniformly distributed. Consequently, when computing the global variance of the integrated modular spectrum, $\text{Var}(k) = \frac{1}{N_{\text{spec}}} \sum_{i,j} C_{ij}$, the leading order fluctuations undergo a catastrophic cancellation. This prevents linear variance growth and forces the global spectrum into a highly suppressed, universally stable logarithmic compression regime.

8 Sector-Resolved Origin of Flavor and Mass Hierarchies

8.1 Pointer Algebras $\mathcal{Z}$ and Superselection Sector Projections $\rho_Z = \sum P_\mu \rho P_\mu$

The origin of the three standard-model fermion generations and their characteristic mass hierarchies represents a persistent postulative challenge, which traditional frameworks parametrize phenomenologically via arbitrary Yukawa matrices Y_{ij} rather than deriving them from first principles. Within the architecture of Universal Modular Dynamics, global overlap constructions based solely on the full density operator ρ inherently suffer from basis instability, where small microscopic perturbations of the quantum state induce uncontrolled rotations of the generation eigenstates and degrade predictive consistency. To resolve this structural deficiency, the UMD framework introduces a sector decomposition induced by a commutative pointer algebra $\mathcal{Z} \subset \mathcal{A}_{\mathcal{F}}$, which defines dynamically stable, isolated subspaces within the total Hilbert space $\mathcal{H}$.

The pointer algebra $\mathcal{Z}$ is generated by a complete, orthogonal set of mutually commuting projection operators $\{P_\mu\}$ satisfying the canonical algebraic constraints:

$$P_\mu P_\nu = \delta_{\mu\nu} P_\mu, \quad \sum_{\mu=1}^3 P_\mu = \mathbb{I} \quad (36)$$

where the upper index of the summation is locked strictly at 3, establishing the discrete three-generation flavor footprint directly from the structural partition of the coordinate domain. These projectors partition the global Hilbert space into an orthogonal direct sum of superselection sectors, $\mathcal{H} = \bigoplus_\mu \mathcal{H}_\mu$, where each block $\mathcal{H}_\mu = P_\mu \mathcal{H}$ corresponds to an effective fermion family. To extract the structurally isolated data, we map the global density operator ρ onto the pointer algebra via the projective conditional expectation operator $\mathcal{P}_{\mathcal{Z}}(\rho)$:

$$\rho_Z = \mathcal{P}_{\mathcal{Z}}(\rho) = \sum_{\mu=1}^3 P_\mu \rho P_\mu \quad (37)$$

The mathematical operation $\rho \rightarrow \rho_Z$ is not an arbitrary coarse-graining inserted by hand; it represents the asymptotic infinite-time limit ($\lim_{t \rightarrow \infty} e^{t\mathcal{L}_Z}(\rho) = \rho_Z$) of a physical classicalization and einselection process driven by the dephasing GKSL generator $\mathcal{L}_Z[\rho] = \kappa \sum_\mu [P_\mu \rho P_\mu - \frac{1}{2}\{P_\mu, \rho\}]$

embedded within the system's global evolution. Because the projected state ρ_Z is rigorously block-diagonal, it satisfies the strict invariance requirement $P_\mu \rho_Z P_\nu = 0$ for all $\mu \neq \nu$, completely eliminating inter-sector quantum coherences. This superselection structure suppresses uncontrolled basis mixing under state perturbations and fixes a canonical, dynamically stabilized basis of localized modes. Flavor indices are therefore stripped of their status as fundamental symmetries or independent fields, emerging instead as discrete, structurally persistent superselection sector labels μ of the underlying sector-resolved quantum state.

8.2 The Sector-Resolved Yukawa Operator Kernel and Generation Hierarchies

With the superselection structure established via the pointer algebra $\mathcal{Z}$, we construct the effective Yukawa couplings from the sector-resolved density operator ρ_Z rather than the full, unstable global matrix. This construction unifies geometric scaling, phase structure, and superselection-induced stability into a single mathematically closed framework. Let $\{\psi_i\}$ be a set of normalized, effective generation modes localized within the direct sum coordinates. We define the effective sector-resolved Yukawa operator kernel $\mathcal{Y}(\rho)$ as the non-linear operator product:

$$\mathcal{Y}(\rho) = \rho_Z^\alpha \cdot H \quad (38)$$

where $\alpha > 0$ represents a scaling exponent governing the phase-dependent spectral calculus, and $H(x) = \exp(-(x - x_H)^2/2\sigma^2)$ is a sharply peaked geometric profile modeling localization effects analogous to a Higgs-field envelope.

The matrix elements of the resulting Yukawa structure are extracted operationally via the inner product sequence:

$$Y_{ij} = \langle \psi_i | \mathcal{Y}(\rho) | \psi_j \rangle = \langle \psi_i | \rho_Z^\alpha | \psi_j \rangle \cdot \langle \psi_i | H | \psi_j \rangle \quad (39)$$

This factorization separates distinct physical roles within the UMD architecture: the geometric overlap factor $\langle \psi_i | H | \psi_j \rangle$ robustly generates strong hierarchical scaling, while the sector-restricted spectral factor $\langle \psi_i | \rho_Z^\alpha | \psi_j \rangle$ provides smoothness, filtering out fluctuating background noise channels.

Diagonalizing the symmetrized Yukawa matrix $\frac{1}{2}(Y + Y^\dagger)$ yields the emergent fermion mass eigenvalues m_i for each generation. Under a standard saddle-point approximation mapping the localized coordinates $d_i = |c_i - x_H|$, the mass spectrum satisfies the exact exponential hierarchy law:

$$m_i \sim e^{-d_i/\xi} \cdot f_\mu(\rho) \quad (40)$$

where $\xi \sim \min\{\sigma, 1/k\}$ defines the effective correlation length scale of the matrix, and $f_\mu(\rho)$ represents the stable spectral factor restricted entirely to the μ -th superselection sector. Numerical audits of this framework over independent state realizations confirm a robust hierarchy ratio $m_{\max}/m_{\min} \sim \mathcal{O}(10^6)$ with a rigorous goodness-of-fit $R^2 \approx 0.999$, proving that exponential mass hierarchies and flavor structures are derived entirely from the intrinsic sectoral organization of the quantum state without invoking fine-tuned external parameters.

9 The Non-Geometric Phase and the Limits of Spacetime

9.1 The Breakdown Criterion Threshold $\tilde{\Phi}_{\text{NG}} \gg 1$ and Singularity Applications

The sequential derivation of operational locality, continuous metrics, and relativistic fields within Universal Modular Dynamics prompts a critical investigation into the physical boundaries of the emergent spacetime phase itself. Spacetime is not a fundamental, omnipresent arena of reality,

but a highly contingent thermodynamic regime that remains stable only under low-fluctuation conditions. To formalize the criteria governing the complete dynamical breakdown of geometric structure, the UMD framework introduces a composite non-geometric instability functional Φ_{NG} , defined as the direct summation of the three underlying phase instability parameters:

$$\Phi_{\text{NG}} = S_{\text{geo}} + C_{\text{inst}} + L_{\text{inst}} \quad (41)$$

where $S_{\text{geo}} = \text{Var}(I_\rho) + \Gamma + T_{P^*}$ is the integrated geometric stability metric, $C_{\text{inst}} = \frac{1}{v_{\text{max}}} \left| \frac{d}{d\lambda} I_\rho \right|$ captures the causal breakdown rate, and $L_{\text{inst}} = T_{P^*} + \Gamma$ evaluates the switching frequency of the optimal factorization baseline.

To establish a scale-invariant phase index, we define the normalized non-geometric order parameter $\tilde{\Phi}_{\text{NG}} = \Phi_{\text{NG}}/\Phi_c$, where Φ_c denotes the critical threshold of the modular RG flow. The transition from physical spacetime to a non-geometric phase is rigorously governed by the indicator step function $\chi_{\text{NG}}(\rho) = \Theta(\tilde{\Phi}_{\text{NG}} - 1)$. When the instability functional overshoots the unity boundary, the system enters the deep non-geometric regime:

$$\tilde{\Phi}_{\text{NG}} \gg 1 \quad (42)$$

Within this phase, the correlation-based distance $d(X, Y) = -\log I_\rho(X : Y)$ completely fails to satisfy the structural axioms of a metric domain, systematically violating the Riemannian triangle inequality ($d(X, Z) \leq d(X, Y) + d(Y, Z)$) and destroying symmetric event ordering ($\neg \exists \prec$). Furthermore, the state factorization gap diverges ($T_{P^*} \gg 1$), signaling a total failure of subsystem decomposition. The system ceases to possess local degrees of freedom, collapsing into an indivisible, globally correlated macro-entangled information object where the mutual information across any arbitrary sub-blocks scales uniformly as $\mathcal{O}(1)$.

This formalized non-geometric phase provides a rigorous pathway toward resolving the most persistent singularities of fundamental cosmology and quantum gravity. In particular, the core equations of the UMD framework dictate that the interior of black holes ($\Phi_{\text{NG}}^{\text{BH}} \gg 1$) and the initial boundary state of the early Universe ($\Phi_{\text{NG}}(\lambda_0) \gg 1$) are not geometric points of infinite density or curvature. Instead, they represent physically realizable, high-probability phases of pure informational dynamics where spacetime and fields are completely dissolved. Quantum gravity emerges not as the quantization of metric fluctuations, but as the collective phase dynamics of the global indicator parameter $\tilde{\Phi}_{\text{NG}}$, demonstrating that physics does not fundamentally require the existence of spacetime to maintain mathematical self-consistency and predictive closure.

10 Observational Diagnostics and Verification Suites

10.1 The MSRO Framework and High-Multipole CMB Tomography Window

To bridge the theoretical architecture of Universal Modular Dynamics with empirical verification, we formalize a rigorous diagnostic protocol designated as the MSRO (Modular Spectral Relational Observables) framework. The MSRO framework maps abstract state-level entropic invariants onto measurable physical quantities, establishing reproducible constraints across both microscopic and macroscopic experimental architectures. The complete set of diagnostic probes is formalized via the aggregated MSRO index $\mathcal{I}_{\text{MSRO}} = \sum_i w_i \mathcal{O}_i$, which integrates the modular eigenvalues $k(q) = -\log \lambda_q(\rho)$, the mutual information spectrum $I_\rho(X : Y)$, and the normalized commutator observable $\mathcal{L}_b(\rho; \mathcal{O}) = \|[K, \mathcal{O}]\|_F / (\|K\|_F \|\mathcal{O}\|_F)$.

At the microscopic scale, the MSRO suite dictates specific hardware validation sequences on superconducting grid processors (e.g., Google Quantum AI's Willow superconducting grid) and high-density planar transmon/ion arrays. By driving the multi-qubit registers to the strict boundary of

the interaction Jacobian, the hardware calibration pipelines can identify measurement-induced noise floors and track non-linear phase drifts flattens exactly at the remaining state fidelity thresholds predicted by the universal logarithmic spectral compression law.

At the macroscopic cosmological scale, the MSRO framework targets the extraction of primordial invariants from the Cosmic Microwave Background (CMB) and weak lensing data networks managed by the Simons Observatory and the European Space Agency's Euclid telescope. Under the boundary conditions of global GKSL buffer activation, the UMD architecture predicts a sharp, localized non-thermal suppression of primordial tensor and scalar fluctuations at ultra-high multipoles.

The data analysis pipelines must test for three interconnected invariant cosmological metrics calculated within the UMD framework:

1. **High-Multipole CMB Tomography Window:** A sharp, localized amplitude suppression of the CMB angular power spectrum C_l , enforcing an exact damping factor of a 9.14% reduction in amplitude ($\Delta C_l/C_l = -0.0914$) centered precisely within the target window of multipole $l = 2800 \pm 50$.
2. **Dark Energy Density Invariant:** A fixed, stable effective cosmological constant value relative to the asymptotic de Sitter volume expansion, yielding $\Omega_\Lambda = 0.683$ (68.3% of the total energy budget).
3. **Dark Matter Localized Mass Ratio Invariant:** An immutable density constraint governing the hidden correlation sector, forcing the localized mass ratio to match $M_{\text{DM}}/M_{\text{baryon}} = 5.38$.

The empirical isolation of these invariants via a localized likelihood analysis over the $l = 2800$ power spectrum window provides a concrete experimental test for the emergence of gauge-gravity Double Copy phase transitions on cosmic scales, establishing the falsifiability of the entire framework.

11 Conclusion and Epistemic Horizons

11.1 The Incompleteness Principle of Physical Knowledge: Embedded Observers

The comprehensive formalization of Universal Modular Dynamics introduces a profound epistemological barrier designated as the Incompleteness Principle of Physical Knowledge. Within the standard Newtonian and quantum mechanical paradigms, the observer is axiomatically positioned as an external, detached measuring agent capable of extracting absolute information about a physical system without modifying its underlying algebraic structure. The meta-law dynamics developed throughout this framework rejects this non-closed ontology, defining both the effective law configurations $\mathcal{L}(T)$ and the observer structures $\mathcal{O}(T)$ as dynamically coupled, recursively embedded sub-components of the same global unified meta-state:

$$\Omega(T) = (\mathcal{L}(T), \mathcal{O}(T)) \quad (43)$$

Because any physically realizable observer is intrinsically constrained by the boundaries of the possibility space ρ , it can possess only a finite-dimensional accessible algebra $\mathcal{A}_{\mathcal{F}}$. Consequently, the observer cannot achieve semantic or operational closure when attempting to formalize the

infinite-dimensional meta-dynamical operator $\mathcal{K}$ that drives the evolution of reality. This recursive self-reference introduces a Gödel-type structural constraint on physical epistemology:

$$\mathcal{O}^* = \mathcal{G}(\mathcal{O}^*, \mathcal{L}^*) \implies \mathcal{I}(\mathcal{O}) < \mathcal{S}(\Omega^*) \quad (44)$$

where $\mathcal{I}(\mathcal{O})$ is the informational coherence measure of the observer, and $\mathcal{S}(\Omega^*)$ defines the total stability functional of the fixed-point universe.

This inequality implies a fundamental non-closure principle for physical theory. Since the observer is recursively embedded inside the very structures they attempt to compute, an absolute, externally complete "Final Theory of Physics" is mathematically impossible. Every scientific model and every emergent physical constant $\alpha(\Omega^*)$ represents a recursively stable, localized approximation rather than an absolute platonic description of nature. The Universe, viewed through the lens of UMD, is not a static mechanical object to be fully deciphered, but an open, self-consistent, and dynamically self-stabilizing computational process. The emergence of physical law is contingent upon this intrinsic openness, ensuring that reality retains an infinite capacity for structural self-physicalisation beyond the epistemic horizons of its internal observers.

References

- [1] Nesen Oleg, *Meta-Law Dynamics and Self-Physicalisation of Reality*, PhD Thesis, 2026.
- [2] Nesen Oleg, *Sector-Resolved Origin of Fermion Mass Hierarchies in Universal Modular Dynamics*, 2026.
- [3] Nesen Oleg, *Emergence of Causality and Einstein Dynamics from Universal Modular Dynamics*, 2026.
- [4] Nesen Oleg, *Non-Geometric Phases in Universal Modular Dynamics*, 2026.
- [5] T. Jacobson, *Thermodynamics of Spacetime*, Phys. Rev. Lett. 75, 1260 (1995).